



## Original Article



## Frequency of Urinary Incontinence and Its Risk Factors in Patients Attending Gynae Outpatient Department (OPD) of Lady Reading Hospital (LRH) Peshawar, Pakistan

Asiya Javed<sup>1</sup>, Sadia Bibi<sup>2</sup> and Alia Firdus<sup>1</sup><sup>1</sup>Type C Hospital Takhte Nasrati, Karak, Pakistan<sup>2</sup>Ujala Medical Complex, Ghorghushti, Pakistan

## ARTICLE INFO

## Keywords:

Women, Urinary incontinence, Factors, Lower Urinary Tract Infection

## How to Cite:

Javed, A., Bibi, S., & Firdus, A. (2025). Frequency of Urinary Incontinence and Its Risk Factors in Patients Attending Gynae Outpatient Department (OPD) of Lady Reading Hospital (LRH) Peshawar, Pakistan: Urinary Risk Factors in Patients Attending Gynae Outpatient Department. *Pakistan Journal of Health Sciences*, 6(3), 126-130. <https://doi.org/10.54393/pjhs.v6i3.1103>

## \*Corresponding Author:

Asiya Javed  
Type C Hospital Takhte Nasrati, Karak, Pakistan  
[javedash101@gmail.com](mailto:javedash101@gmail.com)Received date: 10<sup>th</sup> July, 2024Acceptance date: 16<sup>th</sup> March, 2025Published date: 31<sup>st</sup> March, 2025

## ABSTRACT

Urinary incontinence (UI) is an important health issue acknowledged by the World Health Organization (WHO) that adversely affects the quality of life. **Objectives:** To find out the occurrence of UI in female presenting to the Gynaecology Out-Patient Department of Lady Reading Hospital and also to find out the frequencies of its risk factors for urinary incontinence in women presenting with UI. **Methods:** A descriptive cross-sectional design was used to conduct this study. The sample consisted of 163 women who presented to the Gynaecology Out-Patient Department of Lady Reading Hospital for any complaints were included in the study. **Results:** UI was observed in 15.3% of patients. Moreover, Stress UI was 48%, followed by overflow UI 24%, urge UI 20%, and mixed UI 8%. The leading factors for UI consisted of 3.75% more than three normal vaginal deliveries, 24% chronic cough, 12% history of gynaecological operation, 80% Illiterate, and 56% lower urinary tract infection (LUTI). **Conclusions:** It was concluded that based on the findings of the current study, UI is 15.3% prevalent in Pakistan which is comparable to that of other Asian nations. Furthermore, normal vaginal delivery more than three times and UTI may develop the key risk issues for the development of UI. The magnitude of UI problems may impact on daily lives and habits of the patients.

## INTRODUCTION

The International Incontinence Society defines urinary incontinence (UI) as a social and hygienic issue where involuntary urine leakage is objectively observable [1]. After giving birth, over 29.6% of women reported having urine incontinence [2]. Its prevalence in the Turkish population has been estimated to reach 50% [3]. According to Oman research, 28.25% of the women were between the ages of 50 and 59 years. Among Omani women aged 20 to 60, the point prevalence (per 1000) of UI was 44%. Stress UI (41.6%) is found in the majority of the women. Furthermore, 15.2% of women with UI had mild UI, 50.3% had moderate UI, 33.1% had severe UI, and just 1.3% had extremely intense UI [4]. Similarly, a research study in

Indonesia estimated the prevalence of urinary incontinence at 13.0% overall [5]. Moreover, a study in India analysed that 27.7% of women have urinary incontinence in which urge UI (16.3%) followed by mixed UI (32.7%) and stress UI (51.0%) [6]. Furthermore, a similar study in India showed that the overall prevalence of urinary incontinence in patients presenting to the hospital was about 12% [7]. According to research done in Turkey, 37.2% of these women experienced urine incontinence, however, only 29.3% of them sought medical attention for the issue. Stress-type incontinence was considered the most observable type in 160 women with urine incontinence findings (33.7%), followed by other types (overflow,



continuous urinary incontinence) (13.7%), urge type (20.6%) and mixed type (31.8%) [8]. Forty-one (10.30%) of the women were diagnosed with stress urine incontinence. Tuberculosis and other lung conditions were found to be the most prevalent co-morbidities. Only 4% of the women in the trial developed a UTI that tested positive for culture [9]. Research has shown that more than three normal vaginal births, chronic cough, history of gynaecological procedures, and illiteracy have all been identified as the key risk factors for UI [10]. UI has been shown to have negative effects on sexual relationships, work/school performance, and family life [11]. Among women over 40, UI was described to be significantly higher (69.7%). Out of 198 participants, 179 (90.4%) were from the poor socioeconomic group, followed by 17 participants from the middle class, and 2 (1%) were from the higher socioeconomic group. Increasing parity was linked to incontinence, particularly stress incontinence since 81 patients (40.9%) had grand multipara, which is a substantial risk factor for urine incontinence. Additional risk factors were a history of prolapse (34.8%), constipation (51%), elevated BMI > 25 per kg/m<sup>2</sup> (47%), vaginal birth (85.4%), and chronic respiratory illness (10%) [12]. Numerous pregnancies, UI in the mother and sister, prior UI during pregnancy, the postpartum period, and body mass index were found to be linked to an increased risk of incontinence. Just 40 out of 92 women (43.5%) have sought medical assistance, whereas 92 out of 288 women (31.9%) said that UI negatively affects their quality of life. The kind of UI, the frequency of UI episodes, and the degree of leakage were all associated with a worse quality of life [13]. Age, vaginal birth, and body mass index (BMI) all contributed to an increase in the prevalence of urine incontinence. Stress urinary incontinence (45.2%), urgency urinary incontinence (22.0%), and mixed urinary incontinence (32.8%) were reported by participants with undiagnosed UI. Urinary incontinence was shown to be significantly correlated with constipation and a high body mass index [14]. In women with Stress Urinary Incontinence (SUI), Mid Urethral Sling (MUS) surgery and Pelvic Floor Muscle Training (PFMT) dramatically enhance quality of life. In the surgical intervention arm, we saw 100% symptomatic alleviation, a high rate of Quality of Life (QoL) improvement, and few easily manageable sequelae. Despite improving quality of life, PFMT needs ongoing encouragement because 22% of participants stopped using it. In the absence of medication, SUI patients' quality of life was rather constant [15].

This study aims to find out the occurrence of UI in female presenting to the Gynaecology Out-Patient Department of Lady Reading Hospital and also to find out the frequencies of its risk factors for urinary incontinence in women presenting with UI.

## METHODS

A descriptive cross-sectional approach was applied to conduct this study in the Obstetrics and Gynecology unit of Lady Reading Hospital (LRH), Peshawar. Using Openepi software, an anticipated population Proportion (P) of urinary incontinence in patients presenting to the hospital was about 12% [7], confidence interval of 95% and a 5% margin of error, the calculated sample size was 163 women through non-probability consecutive sampling technique. This study was completed in six months starting from July 1<sup>st</sup> to December 31<sup>st</sup>, 2020. The written approval of the ethical committee of the hospital was granted before the conduct of this study. Along with this, study participants signed written informed consent at the start of the data collection. The quantitative data was collected through a structured questionnaire Performa. The inclusion criteria include all those women who presented to Gynae OPD of LRH for any complaint, having aged between 25 to 65 years, gravidity and parity both greater than one and willing to take part in this research. Similarly, all those women having known carcinoma of the genital tract, pregnancy, obstetric problems, and present for any type of prolapse like uterine prolapse and vaginal prolapse were excluded. SPSS version 23.00 software was used for data analysis. Mean and standard deviation were calculated in descriptive statistics for continuous variables like participants' age, gravidity and BMI. Percentage and frequency were calculated for categorical variables like urinary incontinence, different types of UI and for each risk factor. The inferential statistics include a Chi-square test for an association between the dependent variable (Urinary Incontinence) and independent variables (Selected Demographics). All the results were statistically significant when the p-value was <0.5 at a 95% confidence interval.

## RESULTS

The mean age of the participants in this study was 42.65 ± 7.37 years; moreover, the mean gravida was 4.25 ± 1.04 and the mean BMI was 27.64 ± 2.21 Kg/m<sup>2</sup>. 15.3% of the sample population in the current study have UI; followed by 84.7% of participants who have no UI. Similarly, 48% of the participants have stress UI, followed by 24% overflow UI, 20% Urge UI, and 08% have Mixed UI in the current study (Table 1).

**Table 1:** Urinary Incontinence and Its Sub-Types (n=163)

Sr. no	Variables	Frequency	Percentage
<b>Urinary Incontinence</b>			
1.	Yes	25	15.3%
2.	No	138	84.7%
<b>Urinary Incontinence Sub-Types (n=25)</b>			
1.	Stress UI	12	48%
2.	Overflow UI	06	24%

3.	Urge UI	05	20%
4.	Mixed UI	02	08%

More than three normal vaginal deliveries accounted for 30.64% of UI, followed by 09.67% chronic cough, 04.83% history of gynaecological operations, 32.25% educational status of the participants, and 22.58% lower urinary tract infection respectively. Whereas the percentage of individual risk factors shows that 76% of UI patients have more than three normal deliveries and 24% of UI patients have less than three normal vaginal deliveries. Furthermore, 24% of UI patients have chronic cough and 76% of UI patients have no chronic cough. 12% of UI patients have a history of gynaecological operations and 88% of UI patients have no history of gynaecological operations. 80% of UI patients were illiterate and 20% of UI patients were literate. Similarly, 56% of UI patients have lower urinary tract infections and 44% of UI patients were negative for lower urinary tract infections (Table 2).

**Table 2:** Risk Factor of Urinary Incontinence (n=25)

Sr. no	Risk factors	Frequency (%)	Commuted %
<b>Normal Vaginal Deliveries &gt; 3</b>			
1.	Yes	19 (76%)	30.64%
2.	No	06 (24%)	09.52%
<b>Chronic Cough</b>			
1.	Yes	06 (24%)	09.67%
2.	No	19 (76%)	30.15%
<b>History of Gynecological Operations</b>			
1.	Yes	03 (12%)	04.83%
2.	No	22 (88%)	34.92%
<b>Educational Status</b>			
1.	Illiterate	20 (80%)	32.25%
2.	Educated	05 (20%)	07.93%
<b>Lower Urinary Tract Infection</b>			
1.	Yes	14 (56%)	22.58%
2.	No	11 (44%)	19.04%

UI and selected demographics in the current study were checked for an association through the Chi-square test. It was evident that there was no association between UI and age categories as the p-value ( $0.626 > 0.05$ ) at a 95% confidence interval. Furthermore, UI and gravidity were not associated with each other as the p-value ( $0.930 > 0.05$ ) at a 95% confidence interval. However, there was an association between UI and BMI in the current study as the p-value ( $0.013 < 0.05$ ) at a 95% confidence interval (Table 3).

**Table 3:** Chi-Square test between Urinary Incontinence and Selected Variables

Sr. no	Variables	Urinary Incontinence		p-Value
		Yes	No	
1.	Age Categories			0.626
	25 – 40 Years	12 (16.9%)	58 (83.1%)	
	41 – 65 years	13 (14.1%)	79 (85.9%)	
2.	Gravida			0.930
	1 – 3	06 (15.8%)	32 (84.2%)	
	>3	19 (15.2%)	106 (84.8%)	
3.	BMI (Kg/m <sup>2</sup> )			0.013*
	≤25	00 (0%)	28 (100%)	
	>25	25 (15.3%)	138 (81.5%)	

\*p<0.05 Significance Level and at 95% Confidence Interval

## DISCUSSION

This research reported a 15.3% prevalence of urine incontinence, which was lower than the 25–45% noted in European studies [9]. Garg *et al.*, revealed that 21.8% of Indian women had UI [16]. Because UI is frequently seen as a demeaning disease in Pakistani and other Asian cultures, the decreased incidence in this study may be because patients did not report their experiences with UI. Nonetheless, it is predicted that the number of UI patients will increase more in Asia (22%) than in the United States (18%) and Europe (5%) [17]. It suggests that the iceberg phenomenon and the scope of UI issues in Asian locations are comparable. In the female population of this study, the most prevalent UI was the stress type UI. The stress-type UI was the most common among the study's female participants. Garg *et al.*, and Rashidi *et al.*, also find similar results. There are several potential risk factors for stress UI in women, including decreased estrogen levels during the menopausal period, pelvic floor muscle, nerve, and connective tissue injury that occurs throughout pregnancy to delivery, and a history of gynaecological surgery [16, 18]. These many causes can result in urethral hypermobility, problems in the intrinsic structure of the urethral sphincter, and damage to the urethral supporting tissue (the levator ani, the anterior vaginal wall, and the extrinsic structure of the urethra), all of which can be signs of stress UI [19]. Multiparity and urodynamic parameters are the primary risk factors for the development of UI in Lower Urinary Tract infections (LUTI), according to retrospective research conducted on Japanese female patients. The characteristics of LUTI in urodynamics, maximum free flow rate, maximal urethral closure pressure, and functional profile length, are linked to a higher incidence of UI in female patients [14]. Urinary incontinence risk factors include metabolic syndrome, spontaneous birth, inadequate sanitation, advanced age, and urinary tract infections, according to a different study conducted on Chinese women [20]. In the present study, stress urinary

incontinence was 48%, Overflow urinary incontinence 24%, Urge urinary incontinence 20% and mixed urinary incontinence 8%. Factors leading to urinary incontinence were normal vaginal delivery >3 76%, chronic cough 24%, history of gynaecological operation 12%, Illiterate 80% and lower urinary tract infection 56%. UI impairs social life components, physical activity, sexual relationships, and religious responsibilities. Furthermore, Age, UI intensity, UI kind, frequency of episodes, body weight, psychological variables, and medical treatment patterns are the elements that impact UI patients' quality of life.

## CONCLUSIONS

It was concluded that UI is 15.3% prevalent in Pakistan, comparable to other Asian nations. Furthermore, low educational status (Illiterate) accounted for 32.25% of UI followed by 30.64% normal vaginal delivery >3, 22.58% Lower urinary tract infection, 09.67% Chronic cough, and 04.83% history of gynaecological operations. UTI may become the main risk factor in the development of UI. In addition, normal vaginal delivery more than three times and UTI may develop the key risk issues for the development of UI. The magnitude of UI problems may have an impact on the daily lives and habits of the patients.

## Authors Contribution

Conceptualization: AJ

Methodology: AJ, AF

Formal analysis: SB, AF

Writing review and editing: SB

All authors have read and agreed to the published version of the manuscript

## Conflicts of Interest

All the authors declare no conflict of interest.

## Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

## REFERENCES

- [1] Srisanam M, Buppasiri P, Chongsomchai C, Temtanakitpaisan T. Urinary Incontinence: Women Attending the Gynecology Outpatient Clinic Unaware of Symptoms. *Thai Journal of Obstetrics and Gynaecology*. 2022 Mar; 136-44.
- [2] Fahim F. The Frequency, Types and Risk Factors of Urinary Incontinence in Women Presenting to the Gynaecology Outpatients of a Tertiary Care Hospital. *Journal of Medical Sciences*. 2021;29(04):273-6. doi: 10.52764/jms.21.29.4.13.
- [3] Kershaw VF, Chainrai M, Radley SC. Patient Initiated Follow Up in Obstetrics and Gynaecology: A Systematic Review. *European Journal of Obstetrics and Gynaecology and Reproductive Biology*. 2022 May; 272: 123-9. doi: 10.1016/j.ejogrb.2022.02.181.
- [4] Seshan V, Francis F, Raghavan D, Arulappan J, Hashmi IA, Prince EJ et al. Prevalence of Urinary Incontinence and Its Relationship with Sociodemographic and Obstetrical Variables Among Omani Women. *SAGE Open Nursing*. 2023 May; 9: 23779608231173803. doi: 10.1177/23779608231173803.
- [5] Sheyla Almeida Gomes Braga FD, Vasconcelos Benício CD, Gonçalves Bezerra SM, da Silva A, Quaresma Costa A, Soares dos Santos E et al. Profile of Patients with Urinary Incontinence in A University Hospital Outpatient Clinic. *Revista Estima*. 2021 Jan; 19.
- [6] Raabo SA, Swarray-Deen A, Sefogah PE. Prevalence of Urinary Incontinence and Associated Factors among Patients Attending Gynaecology Clinic at Korle-Bu Teaching Hospital, Accra, Ghana. *Gynecology Reproductive Health*. 2025; 9 (1): 1-5. doi: 10.31579/2578-8965/257.
- [7] Salkapuram SK, Elumalai K, Williams H, Sivannan S, Srinivasan S, Anandakumar S. Prevalence, Risk Factors, and Statistical Analysis of Urinary Incontinence in a Tertiary Care Hospital in India. *Prevalence*. 2024 Nov; 25(4): 286-92. doi: 10.4274/imj.galenos.2024.09514.
- [8] Kılıç M. Incidence and Risk Factors of Urinary Incontinence in Women Visiting Family Health Centers. *Springer Plus*. 2016 Dec; 5: 1-9. doi: 10.1186/s40064-016-2965-z.
- [9] Bora A and Gupta AS. Prevalence and Assessment of the Risk Factors of Stress Urinary Incontinence in Gynaecology Outpatients in A Tertiary Care Centre. *International Journal of Reproduction, Contraception, Obstetrics and Gynaecology*. 2020 Dec; 9(12): 4959. doi: 10.18203/2320-1770.ijrcog20205229.
- [10] Molinet Coll C, Martínez Franco E, Altimira Queral L, Cuadras D, Amat Tardiu L, Parés D. Hormonal Influence in Stress Urinary Incontinence During Pregnancy and Postpartum. *Reproductive Sciences*. 2022 Aug; 29(8): 2190-9. doi: 10.1007/s43032-022-00946-7.
- [11] Zhang RQ, Xia MC, Cui F, Chen JW, Bian XD, Xie HJ et al. Epidemiological Survey of Adult Female Stress Urinary Incontinence. *BioMed Central Women's Health*. 2021 Dec; 21: 1-0. doi: 10.1186/s12905-021-01319-z.
- [12] Mumtaz A, Saif N, Salah-Uddin N, Siddique U, Nouman D, Farooq F. Original Articles Frequency and Associated Risk Factors of Urinary Incontinence and Its Impact On Life of Women. *Journal of Akhtar Saeed Medical and Dental College*. 2020; 2(1): 19-23. doi: 10.51127/JAMDCV02I01OA04.
- [13] Kocaöz S, Talas MS, Atabekoglu CS. Urinary Incontinence among Turkish Women: An Outpatient Study of Prevalence, Risk Factors, and Quality of Life.



- Journal of Wound Ostomy and Continence Nursing. 2012 Jul;39(4):431-9. doi:10.1097/WON.0b013e318259c42b.
- [14] Srisanam M, Buppasiri P, Chongsomchai C, Temtanakitpaisan T. Urinary Incontinence: Women Attending the Gynecology Outpatient Clinic Unaware of Symptoms. Thai Journal of Obstetrics and Gynaecology. 2022 Mar; 136-44.
- [15] Pandey D, Maturi C, Dhakar BP, Jain G, Kyalakond K. Interventions and Quality of Life in Stress Urinary Incontinence. Gynecology and Minimally Invasive Therapy. 2019 Jul; 8(3): 106-12. doi: 10.4103/GMIT.GMIT\_72\_18.
- [16] Garg P, Goyal LD, Goyal S, Verma M. Utility of Opportunistic Screening to Assess the Impact of Urinary Incontinence On Quality of Life and Barriers to Seeking Treatment Among Women Attending a Tertiary Healthcare Centre in North India. BioMed Central Urology. 2024 Mar; 24(1):50. doi: /10.1186/s12894-024-01434-7.
- [17] Ma T, Wan M, Liu G, Zuo X, Yang X, Yang X. Temporal Trends of Inflammatory Bowel Disease Burden in China from 1990 to 2030 with Comparisons to Japan, South Korea, the European Union, the United States of America, and the World. Clinical Epidemiology. 2023 Dec; 583-99. doi: 10.2147/CLEP.S402718.
- [18] Rashidi Fakari F, Hajian S, Darvish S, Alavi Majd H. Explaining Factors Affecting Help-Seeking Behaviors in Women with Urinary Incontinence: A Qualitative Study. BioMed Central Health Services Research. 2021 Dec; 21:1-0. doi: 10.1186/s12913-020-06047-y.
- [19] Escura Sancho S, Ribera Torres L, Castelo-Branco Flores C, Anglès Acedo S. Impact of Urinary Incontinence on Women's Sexuality. Clinical and Experimental Obstetrics and Gynecology. 2022 Feb; 49(2): 049-054. doi: 10.31083/j.ceog4902049.
- [20] Aksoy YE and Yilmaz SD. The Effect of Urinary Incontinence Symptoms On Sexual Functions and Quality of Life in Women. Sexuality and Disability. 2022 Mar 1:1-3. doi: 10.1007/s11195-021-09714-y.