



Original Article



Assessment of the Proximal Femoral Nailing and Dynamic Hip Screws in Intertrochanteric Fractures

Muhammad Rizwan Ali^{1*}, Irfan Ahmad¹, Nauman Naeem Ahmed¹, Muhammad Farhan Shahzad¹, Yasir Umer², Lavisha³ and Shafqat Shahzad⁴

¹Department of Orthopedics Unit 2, Jinnah Hospital, Allama Iqbal Medical College, Lahore, Pakistan

²Ali Medical Centre, Islamabad, Pakistan

³College of Physiotherapy, Jinnah Postgraduate Medical Centre, Karachi, Pakistan

⁴College of Statistical and Actuarial Sciences, University of the Punjab, Lahore, Pakistan

ARTICLE INFO

Keywords:

Intertrochanteric Fracture, Proximal Femoral Nailing, Dynamic Hip Screw, Femur Fracture Management, Orthopedic Surgery

How to Cite:

Ali, M. R., Ahmad, I., Ahmed, N. N., Shahzad, M. F., Umer, Y., Lavisha, ., & Shahzad, S. (2024). Assessment of the Proximal Femoral Nailing and Dynamic Hip Screws in Intertrochanteric Fractures: Femoral Nailing and Hip Screws. Pakistan Journal of Health Sciences, 5(10). <https://doi.org/10.54393/pjhs.v5i10.1892>

*Corresponding Author:

Muhammad Rizwan Ali
Department of Orthopedics Unit 2, Jinnah Hospital,
Allama Iqbal Medical College, Lahore, Pakistan
drizwan70@gmail.com

Received Date: 17th August, 2024

Acceptance Date: 20th October, 2024

Published Date: 31st October, 2024

ABSTRACT

An intertrochanteric fracture is one of three types of proximal femur fractures, occurring in the proximal part of the femur. The other two types are subtrochanteric fractures, which occur below the trochanters, and proximal (or cephalad) fractures of the femoral neck. **Objective:** To assess the differences in hospital stay, blood loss, and operative time between proximal femoral nailing and dynamic hip screws in the management of intertrochanteric fractures of the femur. **Methods:** Quasi-experimental study was conducted in Unit II of the Orthopedic Surgery department at Jinnah Hospital Lahore. It included eighty patients whose demographic profiles (name, age, gender, and BMI) were collected. Using a non-probability consecutive sampling technique, patients were divided into two groups: Group A underwent Dynamic Hip Screw (DHS) surgery, while Group B received Proximal Femoral Nailing (PFN) surgery. Data on operative time, blood loss, and hospital stay were analyzed using SPSS version 21.0, with outcomes compared via independent sample t-test at $p < 0.05$ significance. **Results:** Regarding hospital stay, blood loss, and operating time, DHS and PFN differed significantly. Compared to the DHS group, the PFN group's operating time, blood loss, and length of hospital stay were much lower. **Conclusions:** This study has demonstrated that PFN is a far superior option to DHS for treating femur intertrochanteric fractures. Going forward, suggested PFN rather than DHS for femur fractures of this kind.

INTRODUCTION

An intertrochanteric fracture is one of three types of proximal femur fractures, occurring in the upper portion of the femur. The two other forms of proximal femur fractures are subtrochanteric fractures, which occur distal to or below the trochanters, and proximal or cephalad fractures of the femoral neck [1]. While it's usual to refer to all three types of fractures as just proximal femur fractures, it's crucial to understand the differences between them as each has unique anatomy, prognosis, and management. Literature from the early 1800s showed that intertrochanteric proximal femur fractures were malunited in varus and often healed, resulting in deformity and reduced function related to limp and weak hip abductor

muscles. However, because of coexisting medical conditions and extended incumbency that hindered union, nonoperative treatment of intertrochanteric fractures yielded notable, intolerable morbidity and death. Regardless of the mortality, conservative treatment was ruled unsuitable due to these serious issues [2]. Consequently, decreasing the degree of malunion and potential nonunion of these fractures, as well as the mortality from non-orthopedic comorbid or concurrent medical issues, have been key components of advancement in the treatment of intertrochanteric fractures. Patients can now function better because of these advancements [3]. The elderly population's

concomitant osteoporosis or osteopenia is the other issue linked to intertrochanteric fractures. Programs to make sure these people are evaluated and, if needed, treated to avoid a second fracture have been developed as a result of this problem's recognition. Given the continuous rise in the older population, it's especially critical to prevent the initial fracture [4]. Simple fracture healing is aided by fracture stability, which is the fracture pattern's capacity to withstand weight-bearing deformation. Furthermore, the various sub-types of intertrochanteric fractures have differing rates of sequelae, including mortality and morbidity, as well as postoperative treatment [5]. Currently, surgery is required to treat intertrochanteric fractures. Even if nonsurgical procedures offer adequate recovery rates, surgical methods have supplanted earlier nonsurgical methods such as prolonged traction in bed, prolonged bed rest, or prolonged immobilization in a full-body cast [6]. The nonsurgical management yielded acceptable healing rates, but the resulting unsatisfactory morbidity and death were a result of bad unions that compromised patient function and numerous non-orthopedic problems linked to extended immobility or inactivity. Among these issues were the following complications [7]. Extended immobilization can lead to various complications, including inactivity-related pneumonia and pulmonary complications, pulmonary embolism due to extremity immobility from deep vein thrombosis, pressure sores from prolonged bed rest, and muscular atrophy with decreased range of motion in the lower extremities. Presently, almost all trochanteric fractures are treated with surgery, which is referred to as open reduction and internal fixation, with a few notable exceptions. Almost any trochanteric fracture can be treated with one of the many internal fixation devices that are available. Every instance requires a precise fit between the right device and the particular kind of fracture. It is also recommended to employ the appropriate surgical method as specified by the device developer. As a result, the ideal device is selected following a careful diagnosis of the trochanteric fracture type. The patient's degree of osteoporosis, activity level, and reasonable expectations for the procedure must all be considered when determining the technique's indications and contraindications [8]. At the very least, technically sound preoperative radiographs of the hip are needed to achieve this match between technique and patient. These radiographs should include a true lateral view (cross-table technique). Sometimes the fracture cannot be well defined to allow for precise operative planning. In these situations, a computed tomography (CT) scan, a frog lateral view, or even a reconstructed CT scan, may be required. If there has been severe shortening, then gentle traction during the radiograph will help characterize the fracture (traction view) [9].

METHODS

A quasi-experimental study was conducted at Unit II of the Orthopedic Surgery Department at Jinnah Hospital in

Lahore over six months, following the approval of the study synopsis. A total of eighty patients were included, with the sample size calculated based on a 95% confidence level and 80% power of the test, using the mean operative times of 59.16 ± 16.92 minutes for Proximal Femoral Nailing (PFN) and 87.35 ± 21.29 minutes for Dynamic Hip Screw (DHS) in intertrochanteric femur fractures [4]. Informed consent was obtained from all participants, and demographic profiles (name, age, gender, and BMI) were collected. Using a non-probability consecutive sampling technique, patients were divided into two groups: Group A underwent DHS surgery, while Group B received PFN surgery. The study included patients aged 20 to 70 years of either gender who presented with a femoral intertrochanteric fracture within 7 days of injury. Exclusion criteria included osteoporosis, osteoarthritis, osteomalacia, and a positive rheumatoid arthritis factor (RA > 14 IU/ml), as well as patients with comminuted fractures, multiple fractures, or open fractures showing signs of infection and debris (e.g., pus discharge) during clinical examination. All surgeries were performed under general anesthesia by a single surgical team to ensure that patients were completely unconscious and pain-free, allowing for optimal muscle relaxation and access to the surgical site. Following anesthesia, the skin was incised, and the operative time was recorded. Total blood loss was measured during the surgery, and once the skin was closed, the final operative time was noted. Patients were then transferred to post-surgical wards, where they were monitored until discharge, and the total hospital stay was documented according to operational definitions. Data were entered and analyzed using SPSS version 21.0. Quantitative variables such as age, BMI, operative time, blood loss, and hospital stay were presented as mean and standard deviation, while qualitative variables like gender and anatomical side were reported as frequency and percentage. Outcomes between the two groups were compared using an independent sample t-test, with a significance level set at $p < 0.05$. After surgery, patients were monitored in post-surgical wards until discharge, and the total hospital stay was recorded.

RESULTS

The findings of the study indicated that for patients under 50 years old, the average operating time was 55.71 ± 3.28 minutes in the PFN group and 74.54 ± 7.75 minutes in the DHS group. For patients over 50 years old, the average operating time was 53.11 ± 2.80 minutes in the PFN group and 79.29 ± 7.33 minutes in the DHS group. For patients under 50 years old, the mean blood loss was 247.16 ± 32.44 ml in the PFN group and 445.08 ± 28.67 ml in the DHS group. For patients over 50 years old, the mean blood loss was

247.33 ± 34.14 ml in the PFN group and 444.50 ± 38.78 ml in the DHS group. The mean hospital stays for patients under 50 years old were 8.71 ± 2.65 days in the PFN group and 10.69 ± 2.09 days in the DHS group. For patients over 50 years old, the mean hospital stays were 8.78 ± 3.07 days in the PFN group and 10.14 ± 1.75 days in the DHS group. Except for the hospital stay for patients over 50, or p-value < 0.05, there was a statistically significant difference seen between the study groups for operative time, blood loss, and hospital stays stratified by age (Table 1).

Table 1: Comparison Between Study Groups Categorized by Age, In Terms of Operative Time, Blood Loss, And Hospital Stay

Study Group	Age (Years)	Study Groups (Mean ± SD)		p-Value
		PFN	DHS	
Operative Time	≤50	55.71 ± 3.28	74.54 ± 7.75	0.001*
	>50	53.11 ± 2.80	79.29 ± 7.33	0.001*
Blood Loss	≤50	247.16 ± 32.44	445.08 ± 28.67	0.001*
	>50	247.33 ± 34.14	444.50 ± 38.78	0.001*
Hospital Stays	≤50	8.71 ± 2.65	10.69 ± 2.09	0.003*
	>50	8.78 ± 3.07	10.14 ± 1.75	0.250

The study's findings revealed that, for male patients, the average operating time was 54.86 ± 3.48 minutes in the PFN group and 76.12 ± 8.61 minutes in the DHS group; for female patients, the average operating time was 55.44 ± 3.22 minutes in the PFN group and 76.33 ± 6.67 minutes in the DHS group. For male patients, the average blood loss was 246.45 ± 36.49 ml in the PFN group and 442.92 ± 29.53 ml in the DHS group. For female patients, the average blood loss was 248.11 ± 27.54 ml in the PFN group and 448.13 ± 26.92 ml in the DHS group. For male patients, the average hospital stays were 8.95 ± 2.72 days in the PFN group and 10.52 ± 1.89 days in the DHS group. For female patients, the average hospital stays were 8.44 ± 2.75 days in the PFN group and 10.47 ± 2.17 days in the DHS group. With operating time, blood loss, and hospital stays stratified by gender, a statistically significant difference (p-value < 0.05) was seen between the study groups (Table 2).

Table 2: Comparison between Study Groups Categorized by Gender, In Terms of Operative Time, Blood Loss, and Hospital Stay

Study Group	Gender	Study Groups (Mean ± SD)		p-Value
		PFN	DHS	
Operative Time	Male	54.86 ± 3.48	76.12 ± 8.61	0.001*
	Female	55.44 ± 3.22	76.33 ± 6.67	0.001*
Blood Loss	Male	246.45 ± 36.49	442.92 ± 29.53	0.001*
	Female	248.11 ± 27.54	448.13 ± 26.92	0.001*
Hospital Stays	Male	8.95 ± 2.72	10.52 ± 1.89	0.003*
	Female	8.44 ± 2.75	10.47 ± 2.17	0.025*

Except for hospital stays with the right anatomical side, or p-value < 0.05, the study's results indicated a statistically significant difference between the study groups in terms of operational time, blood loss, and hospital stay stratified by

anatomical side (Table 3).

Table 3: Comparison between Study Groups Categorized By Anatomical Side, In Terms Of Operative Time, Blood Loss, and Hospital Stay

Study Group	Anatomical Side	Study Groups (Mean ± SD)		p-Value
		PFN	DHS	
Operative Time	Left	55.77 ± 3.22	77.73 ± 7.97	0.001*
	Right	54.33 ± 3.39	73.36 ± 7.02	0.001*
Blood Loss	Left	252.18 ± 32.83	439.88 ± 27.52	0.001*
	Right	241.11 ± 31.66	454.14 ± 28.48	0.001*
Hospital Stays	Left	8.91 ± 2.84	10.69 ± 2.03	0.015*
	Right	8.50 ± 2.59	10.14 ± 1.87	0.055

DISCUSSION

The aging of contemporary human populations has led to a notable increase in the occurrence of intertrochanteric fractures. The DHS is now a routine implant used to treat these fractures and is frequently utilized in extramedullary fixation. Two tools that are frequently utilized in intramedullary fixing are PFN and Gamma Nails. Despite reports of PFN and DHS's effects in the treatment of intertrochanteric fractures, the findings and interpretations vary [10]. The PFN group had an average operative time of 55.13 ± 3.34 minutes, while the DHS group had an average of 76.20 ± 7.85 minutes. The PFN group had an average blood loss of 247.20 ± 32.38 ml, while the DHS group had an average of 444.88 ± 28.34 ml. The hospital stay of the PFN group was 8.73 ± 2.71 days, while the DHS group had an average hospital stay of 10.50 ± 1.97 days. In terms of operation time, blood loss, and hospital stay, the PFN group outperformed the DHS group statistically. According to a study by Zhang K et al., PFN may be a more effective treatment for intertrochanteric fractures than DHS. The PFN group's operating time was noticeably shorter [11]. According to Kumar P et al., research, PFN is superior to DHS in type II intertrochanteric fractures in terms of fewer blood losses, shorter surgical times, earlier weight bearing and mobilization, shorter hospital stays, lower infection risks, and fewer sequelae [6]. Patients who received PFN had lower intraoperative blood loss (73ml), shorter surgery times (91min), and were able to mobilize earlier than those who underwent DHS, which had higher intraoperative blood loss (159 ml) and longer surgery times (105 min) [12-14]. According to another study by Shiraz S et al., and Yan M et al., in the case of DHS, the average hospital stay was 12.4 days (8-14 days), whereas in the case of PFN, it was 7.8 days (4-12 days) (P=0.001). The DHS group took an average of 12 weeks to return to their pre-injury walking abilities, while the PFN group took just 8 weeks (P=0.03) [15, 16]. Yu W et al., conducted a further investigation which revealed that, in contrast to the PFN Antirotation device, the DHS device may not be the best implant for stable intertrochanteric femur fractures. The difference remained over time, reaching 6.4% and 13.4% at the most recent follow-up (P<0.05). Postoperative HHS after 12, 15, 18, 21, 24, 36, and 48 months after surgery, as well as during

the ultimate follow-up, varied statistically. There were no discernible statistical variations in medical problems between the two groups. [17, 18]. Randomized post-operative rehabilitation research by Pajarinen *et al.*, comparing DHS and PFN treatment for peritrochanteric femoral fractures revealed that PFN may enable a quicker recovery of walking ability following surgery than DHS [19]. A randomized trial found that the average operating time for patients treated with PFN was 55 minutes, while the average operating time for patients treated with DHS was 87 minutes. An additional experiment revealed that the average operating time was greater for DHS when compared with PFN. The blood loss was nearly double for DHS than PFN and the duration of hospital stay was also less in PFN [20]. In Type II intertrochanteric femur fractures, PFN is superior to DHS in terms of less blood loss, shorter surgical times, shorter stays in the hospital, lower risk of infection, and other sequelae [21].

CONCLUSIONS

This study has demonstrated that PFN is a far superior option to DHS for the treatment of femur intertrochanteric fractures. Going forward, we shall suggest PFN rather than DHS for femur fractures of this kind.

Acknowledgement

The authors would like to acknowledge the Medical Affairs Department of Getz Pharma for their technical support and assistance in the publication process.

Authors Contribution

Conceptualization: MRA

Methodology: NNA, MFS

Formal analysis: SS

Writing, review and editing: MRA, IA, YU, L, SS

All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Chang SM, Hou ZY, Hu SJ, Du SC. Intertrochanteric femur fracture treatment in Asia: what we know and what the world can learn. *Orthopedic Clinics*. 2020 Apr; 51(2): 189-205. doi: 10.1016/j.jocl.2019.11.011.
- [2] Davis TR, Sher JL, Horsman A, Simpson M, Porter BB, Checketts RG. Intertrochanteric femoral fractures. Mechanical failure after internal fixation. *The Journal of Bone and Joint Surgery British Volume*. 1990 Jan; 72(1): 26-31. doi:10.1302/0301620X.72B1.2298790.
- [3] Haidukewych GJ. Intertrochanteric fractures: ten tips to improve results. *The Journal of Bone and Joint Surgery*. 2009 Mar; 91(3): 712-9.
- [4] Muazzam MA and Ali A. Prevalence of Intertrochanteric Fractures in Young Population with Hip Fractures Presented in Ghurki Trust Teaching Hospital Lahore. *Pakistan Journal of Medical and Health Sciences*. 2022 Aug; 16(06): 727-. doi: 10.5335/0/pjmhs22166727.
- [5] Martinho T and Stoffel K. Treatment of intertrochanteric femur fractures with hip arthroplasty in older patients: a narrative review of indications and outcomes. *Medicina*. 2021 Jul; 57(8): 763. doi: 10.3390/medicina57080763.
- [6] Kumar P, Rajnish RK, Sharma S, Dhillon MS. Proximal femoral nailing is superior to hemiarthroplasty in AO/OTA A2 and A3 intertrochanteric femur fractures in the elderly: a systematic literature review and meta-analysis. *International Orthopaedics*. 2020 Apr; 44: 62333. doi: 10.1007/s00264-019-04351-9.
- [7] Sadeghi C, Prentice HA, Okike KM, Paxton EW. Treatment of intertrochanteric femur fractures with long versus short cephalomedullary nails. *The Permanente Journal*. 2020; 24. doi: 10.7812/TPP/19.229.
- [8] Yıldırım C, Muratoğlu OG, Turan K, Ergün T, Mısırlı A, Aydın M. The intra- and interobserver reliability of five commonly used intertrochanteric femur fracture classification systems. *Joint Diseases and Related Surgery*. 2022 Apr; 33(1): 187. doi: 10.52312/jdrs.2022.498.
- [9] Mallya S, Kamath SU, Annappa R, Nazareth NE, Kamath K, Tyagi P. The Results of Unstable Intertrochanteric Femur Fracture Treated with Proximal Femoral Nail Antirotation-2 with respect to Different Greater Trochanteric Entry Points. *Advances in Orthopedics*. 2020 Mar; 2020(1): 2834816. doi: 10.1155/2020/2834816.
- [10] Wu K, Xu Y, Zhang L, Zhang Y, Xu W, Chu J *et al.* Which implant is better for beginners to learn to treat geriatric intertrochanteric femur fractures: A randomised controlled trial of surgeons, metalwork, and patients. *Journal of Orthopaedic Translation*. 2020 Mar; 21: 18-23. doi: 10.1016/j.jot.2019.11.003.
- [11] Zhang K, Zhang S, Yang J, Dong W, Wang S, Cheng Y *et al.* Proximal femoral nail vs. dynamic hip screw in treatment of intertrochanteric fractures: a meta-analysis. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research*. 2014 Sep; 20: 1628. doi: 10.12659/MSM.890962.
- [12] Jonnes C, Shishir SM, Najimudeen S. Type II intertrochanteric fractures: proximal femoral nailing (PFN) versus dynamic hip screw (DHS). *Archives of Bone and Joint Surgery*. 2016 Jan; 4(1): 23.

- [13] Li XP, Zhang P, Zhu SW, Yang MH, Wu XB, Jiang XY. All-cause mortality risk in aged femoral intertrochanteric fracture patients. *Journal of Orthopaedic Surgery and Research*. 2021 Dec; 16: 1-8. doi: 10.1186/s13018-021-02874-9.
- [14] Marmor M, Guenther G, Rezaei A, Saam M, Matityahu A. Reporting on quality of reduction and fixation of intertrochanteric fractures-A systematic review. *Injury*. 2021 Mar; 52(3): 324-9. doi: 10.1016/j.injury.2021.02.014.
- [15] Shiraz S, Shujaiddin M, Hasan K, Elramadi A, Ahmed G. Comparison of dynamic hip screw and proximal femoral nailing techniques in stable intertrochanteric fractures. *Cureus*. 2023 Jan; 15(1). doi:10.7759/cureus.33366.
- [16] Yan M, Kuang L, Ni J, Ding M, Wang J, Huang J *et al*. Use of a double reverse traction repositrator versus a traction table for the treatment of intertrochanteric femur fractures: a comparative study. *Orthopaedic Surgery*. 2021 Jun; 13(4): 1254-61. doi: 10.1111/os.12956
- [17] Pitts CC, Montgomery TP, Hess MC, Smith WR, Quade JH. Reamed Versus Unreamed Intertrochanteric Femur Fractures, Is It Time?. *Journal of orthopaedic trauma*. 2020 May; 34(5):2527. doi: 10.1097/BOT.0000000000001687.
- [18] Yu W, Zhang X, Zhu X, Yu Z, Xu Y, Zha G *et al*. Proximal femoral nails anti-rotation versus dynamic hip screws for treatment of stable intertrochanteric femur fractures: an outcome analyses with a minimum 4 years of follow-up. *BioMed Central Musculoskeletal Disorders*. 2016 Dec; 17: 1-6. doi: 10.1186/s12891-016-1079-7.
- [19] Pajarinen J, Lindahl J, Michelsson O, Savolainen V, Hirvensalo E. Pertrochanteric femoral fractures treated with a dynamic hip screw or a proximal femoral nail: a randomised study comparing post-operative rehabilitation. *The Journal of Bone and Joint Surgery British Volume*. 2005 Jan; 87(1):76-81. doi: 10.1302/0301-620X.87B1.15249.
- [20] Karayiannis P and James A. The impact of cerclage cabling on unstable intertrochanteric and subtrochanteric femoral fractures: a retrospective review of 465 patients. *European Journal of Trauma and Emergency Surgery*. 2020 Oct; 46: 969-75. doi:10.1007/s00068-018-01071-4.
- [21] Kong M, Tang Y, Tong F, Guo H, Zhang XL, Zhou L *et al*. The analgesic efficacy of pericapsular nerve group block in patients with intertrochanteric femur fracture: A Randomized Controlled Trial. *PLOS One*. 2022 Oct; 17(10): e0275793. doi: 10.1371/journal.pone.0275793.