



Original Article



Efficacy of 0.25% Bupivacaine Alone vs. with Dexmedetomidine for Ultrasound-Guided Supraclavicular Block in Upper Limb Surgery

Sara Sabir¹, Rehan Hassan Khan Niazi², Shahzad Imran¹, Hafiz Faheem Asghar³, Shumaila Ashfaq¹ and Aamna Muzaffar⁴¹Department of Anesthesia, Islam Medical and Dental College, Sialkot, Pakistan²Department of Anesthesia, Ameer Ud Din Medical College, Lahore, Pakistan³Department of Anesthesia, Islam Medical College, Gujranwala, Pakistan⁴Department of Anesthesia, Indus Hospital, Lahore, Pakistan

ARTICLE INFO

Keywords:

Dexmedetomidine, Analgesia, Bupivacaine, Supraclavicular and Sensory Block

How to Cite:

Sabir, S., Niazi, R. H. K., Imran, S., Asghar, H. F., Ashfaq, S., & Muzaffar, A. (2025). Efficacy of 0.25% Bupivacaine Alone vs. with Dexmedetomidine for Ultrasound-Guided Supraclavicular Block in Upper Limb Surgery: 0.25% Bupivacaine with Dexmedetomidine for Ultrasound-Guided Supraclavicular Block. *Pakistan Journal of Health Sciences*, 6(5), 99-103. <https://doi.org/10.54393/pjhs.v6i5.2629>

*Corresponding Author:

Sara Sabir
Department of Anesthesia, Islam Medical and Dental College, Sialkot, Pakistan
sara.92816@gmail.comReceived Date: 3rd December, 2024Revised Date: 8th May, 2025Acceptance Date: 19th May, 2025Published Date: 31st May, 2025

ABSTRACT

The pain after upper limb surgeries can be intense. The brachial plexus block is an effective and commonly used method to manage this pain, reduce opioid use, and facilitate smoother recovery. **Objective:** To see the effect of combining bupivacaine with dexmedetomidine in upper limb surgeries. **Methods:** Quasi-experimental research was carried out at Islam Medical College, Sialkot. 100 patients undergoing upper limb surgeries who were aged between 25 to 65 years. All the patients who were comprised allergies to study drugs, severe organ impairment, coagulopathy, neurological disorders affecting pain perception, and pregnancy were excluded and divided to receive either 0.25% bupivacaine alone or with dexmedetomidine. The outcome variables included pain, analgesia duration, sensory/motor block onset, and 24-hour analgesic consumption. Data were analyzed by SPSS version 23.0. The comparison of quantitative data was done by an independent sample t-test and chi-square test for insightful comparisons between qualitative variables, with the significance level at p-value < 0.05. **Results:** The average age of patients was 40.1 ± 11.5 and 39.8 ± 10.5 years in Group I and Group II, respectively. Male were more in both groups. Group II showed significantly longer analgesia duration (12.7 vs. 5.3 hours), faster sensory/motor block onset, and lower analgesic consumption ($p < 0.001$). There were no significant differences in adverse effects. **Conclusions:** It was concluded that combining dexmedetomidine with bupivacaine significantly extended analgesia, quickened sensory and motor block onset with a reduction in overall consumption of analgesia. Although sedation was more frequent, no major adverse events were observed.

INTRODUCTION

The pain after upper limb surgeries, particularly total shoulder arthroplasty, can be intense. The inter-scalene brachial plexus block is an effective and commonly used method to manage this pain, reduce opioid use, and facilitate smoother recovery [1]. However, the pain relief from a single injection of the anesthetic is usually short-term, even when additional medications are added. Continuous brachial plexus blocks can offer longer-lasting pain relief, but they need specialized expertise and more resources to perform [2]. Supraclavicular block has the

potential to provide the optimal analgesia under regional anesthetic for upper limb treatment when this is the shoulder, arm, elbow, or hand region. This block provides clavicular brachial plexus obstruction and is a reliable and efficient upper extremity anesthesia [3]. Local anesthetics are drugs that are used in regional anesthesia; an example is the common local anesthetic bupivacaine and the supraclavicular block. Bupivacaine is a long-acting amide-type local anaesthetic and the most of the time is specifically used for upper limb regional anesthesia



because of its potent and prolonged blockade of sensory and motor pathways [4]. Bupivacaine offers extended postoperative pain relief by obstructing voltage-gated sodium channels, effectively interrupting nerve signal transmission. Its considerable lipid solubility and strong protein affinity contribute to its long-lasting effects [5]. In the regional anesthesia procedures, like supraclavicular brachial plexus blocks, bupivacaine can uniquely target at lower doses while preserving motor function and the pain fibers. Its gradual dissociation offers prolonged pain relief, reduces the need for opioids, and enhances patient comfort, thus ideal during long surgeries and continuous nerve blocks [6]. To enhance postoperative and regional analgesic potential related to anesthesia there has been research done on various potential adjuvants such as opioids, clonidine and dexmedetomidine [7]. Dexmedetomidine is an α_2 -adrenoceptor agonist and has potential for use as an adjuvant in regional analgesia. The mechanism is that it prevents norepinephrine from being released, thus reducing sympathetic nervous system activity and increasing analgesia [8]. It is also associated with additional benefits as sedation and anxiolysis without significant respiratory depression, which comes very handy in the immediate postoperative era. Dexmedetomidine with bupivacaine has been applied in several techniques of regional anesthesia, and studies have shown mixed results in terms of better block quality, post-operative analgesia and decreased opioid consumption [9, 10]. Unprecedented revolution in the practice of delivering regional blocks with better precision and safety in locating the nerve by Ultrasound-guided (USG) technique [11]. Since it remains in front of the intended target tissue and the anatomical structures, the chances of complications such as nerve damage or accidental vascular puncture are reduced when positioning a needle under ultrasound guidance. Rather, postoperative analgesia is based on an element of reliability and reproducibility that may be improved with the precision of the USG technique [12, 13].

This study aims to compare the effect of combining bupivacaine with dexmedetomidine on the analgesia effect after post-operative recovery time in upper limb surgeries.

METHODS

A quasi-experimental research was conducted at Islam Medical College, Sialkot, from July to November 2024, after ethical approval (Ref: 900/IMCS/ERC/000103). Participants were recruited using a convenience sampling method. Eligibility criteria of participants include patients undergoing upper limb surgeries who were aged between 25 to 65 years and who fall in ASA criteria I and II. All the patients who were comprised allergies to study drugs, severe organ impairment, coagulopathy, neurological disorders affecting pain perception, and pregnancy were

excluded. The Open Epi software is used for sample size calculation by using duration of analgesia in the bupivacaine with dexmedetomidine Group was 722 ± 88.45 min and 210 ± 35.88 in Bupivacaine Alone Group, by taking 80% power of test, 5% margin of Error and 10% drop out rate is 100 (50 in each group) [14]. A written informed consent was taken. A total of 100 participants were enrolled to effectively assess the duration of analgesia and were equally divided into two groups of 50 each. Group I received 20 ml of 0.25% bupivacaine, while Group II received 20 ml of 0.25% bupivacaine combined with 50 μ g of dexmedetomidine for enhanced analgesic effect. Upon arrival in the preoperative area, baseline vital signs and pain scores were assessed by the Visual Analogue Scale (VAS). The total score of VAS was 10, where 0 indicates no pain and 10 shows the worst pain. An 18-gauge intravenous line was secured in each participant, ensuring proper preparation for the procedure. This structured approach reflects our commitment to delivering effective and high-quality pain management. VAS is a widely used tool for pain assessment, demonstrating a reliability of 0.95 [15]. Patients were monitored and pre-medicated with IV midazolam (1 mg). Using a high-frequency linear ultrasound probe, the brachial plexus was visualized, and a 22-gauge needle was used to administer the anesthetic solution according to the randomized group, ensuring safety with incremental injections. The study measured the duration of postoperative analgesia. It also examined the onset of sensory and motor blocks, along with total analgesic consumption within the first 24 hours. In response to pain complaints, rescue analgesia was administered as nalbuphine (0.1 mg/kg). Adverse events, including hypotension, bradycardia, nausea, vomiting, and sedation, were monitored closely. Data analysis was conducted using SPSS version 23.0. Normality of the data using the Shapiro-Wilk test before executing the t-tests. The quantitative variables, including age, VAS score, post op analgesia effect (duration in minutes), and the onset times for sensory and motor blockade through independent sample t-tests. For categorical variables, we utilized the chi-square test for insightful comparisons with the significance level at p -value < 0.05 .

RESULTS

The patients' mean age was 40.1 ± 11.5 and 39.8 ± 10.5 years in Group I and Group II, respectively, with no significant difference ($p=0.82$). Gender distribution was similar, with males comprising 62% in Group I and 52% in Group II, while females made up 38% and 48%, respectively ($p=0.68$). ASA physical status classification was also comparable, with 44% of patients in Group I and 48% in Group II categorized as ASA I, and 56% and 52% as ASA II ($p=0.65$). The mean VAS score at baselines was 6.5 ± 1.2 and 6.7 ± 1.1 in Group I and Group II ($p=0.58$) (Table 1).

Table 1: Clinical Parameters of Patients (n=100)

Characteristics		Group I	Group II	p-value
Age		40.1 ± 11.5	39.8 ± 10.5	0.89
Gender	Male	31 (62%)	26 (52%)	0.68
	Female	19 (38%)	24 (48%)	
ASA Physical Status	I	22 (44%)	24 (48%)	0.54
	II	28 (56%)	26 (52%)	
Baseline Pain Score (VAS)		6.5 ± 1.2	6.7 ± 1.1	0.38


As compared to bupivacaine alone (5.3 ± 1.4 hours), there was a statistically significant prolonged duration of analgesia with (12.7 ± 2.1 hours) addition of dexmedetomidine to bupivacaine ($p < 0.001$). Times of sensory and motor blockade onset were faster in Group II (Bupivacaine + Dexmedetomidine) than in Group I (Bupivacaine alone) at 8.3 ± 1.8 minutes and 12.3 ± 2.6 minutes ($p = 0.002$ and 0.003 , respectively) and total analgesic (40.7 ± 15.8 mg vs 75.4 ± 20.2 mg, $p < 0.001$) requirement was significantly lower among them (Table 2).

Table 2: Comparison of Analgesia Duration and Onset Time, and Total Analgesic Consumption among Study Groups

Characteristics	Group I	Group II	p-value	Test Used
Analgesia Duration				
Analgesia Duration in Hours,	5.3 ± 1.4	12.7 ± 2.1	<0.001	Independent t-test
Onset Time and Total Analgesic Consumption				
Onset of Sensory Block (min)	10.4 ± 2.3	8.3 ± 1.8	0.00*	Independent t-test
Onset of Motor Block (min)	15.2 ± 3.0	12.3 ± 2.6	0.00*	Independent t-test
Total Analgesic Consumption (mg)	75.4 ± 20.2	40.7 ± 15.8	0.000*	Independent t-test

The dextromethorphan and bupivacaine combination produced a significantly greater degree of sedation (16% vs. 0%, $p = 0.01$) without increasing the rate of any other adverse effects, including bradycardia (8% vs. 2%, $p = 0.36$), hypotension (10% vs. 4%, $p = 0.24$), and nausea (4% vs. 6%, $p = 0.65$). While Group II experienced significantly more sedation, the other adverse events were similar (Figure 1).

Adverse event

**Figure 1:** Comparison of Adverse Events in Both Drugs

DISCUSSION

This study evaluates the efficacy of dexmedetomidine as an adjuvant to 0.5% bupivacaine versus bupivacaine alone in spinal anesthesia, aiming to enhance analgesic onset, duration, and quality across upper limb surgery. Literature found that adding dexmedetomidine to bupivacaine

enhances postoperative analgesia, reduces rescue analgesic use, and improves patient satisfaction without increasing severe side effects. It prolongs the duration of sensory evaluation, remains hemodynamically safe, and lowers the postoperative shivering incidence [16]. These findings align with results reported in similar studies, which show that additional trials have further enhanced the benefits of dexmedetomidine supplementation to local anesthetics, thereby improving regional anesthesia, as discussed in the current study. The block provided by bupivacaine combined with dexmedetomidine lasted appreciably longer than bupivacaine alone, reducing postoperative pain and the need for additional analgesics. Furthermore, patients receiving the combined technique reported superior fulfillment with their postoperative pain management [17]. Moreover, another study indicated that incorporating dexmedetomidine with bupivacaine in spinal anesthesia enhances pain relief without raising the risk of adverse effects. The findings indicated that combining dexmedetomidine with bupivacaine enhances postoperative analgesia and accelerates the onset of both motor and sensory nerve block, as observed in the current study. [18]. It is suggested that Dexmedetomidine's contribution to the prolonged instances of analgesia is towards blocking nerve conduction, thus reducing pain signals. Previously conducted studies of brachial plexus blockade revealed that incorporation of Dexmedetomidine with Bupivacaine significantly extended analgesic effects. That our results support these previous findings has been attested, as the combination of Dexmedetomidine and Bupivacaine exhibits significantly longer duration and stronger analgesic effect than Bupivacaine on its own [19]. Dexmedetomidine significantly increased the effects of bupivacaine. There was an earlier onset of time to sensory and time to motor nerve block for Bupivacaine + Dexmedetomidine than Bupivacaine alone. In a like manner, another trial documented a much faster appearance of the sensory and motor blockade and longer analgesia duration when Bupivacaine was combined with Dexmedetomidine [14]. The current study revealed no adverse effects in both study groups, corroborating the safety of both treatments. Although differences in individual side effects were observed, hypotension and bradycardia were more common in bupivacaine plus dexmedetomidine, but were not significant. These findings correspond with previous reports showing that administering dexamethasone shortens the onset time and duration of sensory and motor blockade. There was a faster onset of sensory and motor blocks significantly compared to the control group, with minimal side effects, which confirms the role of dexmedetomidine as a safe and useful adjunct for bupivacaine in regional anesthesia. [20]. Current results showed that bupivacaine when used with dexmedetomidine provided effective analgesia without the need to increase the rate of adverse effects such as

bradycardia (8% vs. 2%, $p=0.36$), hypotension (10% vs. 4%, $p=0.24$) and nausea (There were no major events of high-grade toxicity in neither group, and safety profiles were similar, but not different ($p>0.05$), without significant adverse events being reported. These findings were confirmed by Sane et al., Intriguingly, no serious complications were reported in either group, which implies that adding dexamethasone did not increase the risk of complications[21].

CONCLUSIONS

The combination of dexmedetomidine to bupivacaine significantly enhanced its efficacy. Bupivacaine combined with Dexmedetomidine exhibited a faster onset of both sensory and motor block as compared to Bupivacaine alone. Furthermore, our findings demonstrated that the combination provided effective analgesia with no reported side effects.

Authors Contribution

Conceptualization: SS

Methodology: SS, RHKN, SI, HFA

Formal analysis: RHKN, AM

Writing review and editing: SI, SA

All authors have read and agreed to the published version of the manuscript

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Lehmann LJ, Loosen G, Weiss C, Schmittner MD. Interscalene Plexus Block Versus General Anesthesia for Shoulder Surgery: A Randomized Controlled Study. *European Journal of Orthopedic Surgery and Traumatology*. 2015Feb;25:255-61. doi:10.1007/s00590-014-1483-3.
- [2] Singhal A, Taksande K, Singhal Jr A. Continuous Catheter Techniques Versus Single-Injection Nerve Blocks: A Comprehensive Review of Postoperative Pain Management Strategies. *Cureus*. 2024Sep;16(9). doi:10.7759/cureus.70040.
- [3] D'Souza RS and Johnson RL. Supraclavicular Block. 2023 Jul.
- [4] Riaz MA, Qamar SB, Siddiqi A, Sardar S, Tariq W, Shah SQ, Habib MF. Comparison of Duration of Action in Spinal Anesthesia of 0.5% Bupivacaine and 0.5% Bupivacaine with Dexmedetomidine: Spinal Anesthesia: Bupivacaine vs. Bupivacaine with Dexmedetomidine. *Journal of Health and Rehabilitation Research*. 2024Sep;4(3):1-5. doi:10.61919/jhrr.v4i3.1459.
- [5] Paganelli MA and Popescu GK. Actions of Bupivacaine, A Widely Used Local Anesthetic, On NMDA Receptor Responses. *Journal of Neuroscience*. 2015Jan;35(2):831-42. doi:10.1523/JNEUROSCI.3578-14.2015.
- [6] Hussain N, Costache I, Kumar N, Essandoh M, Weaver T, Wong P et al. Is Supraclavicular Block as Good as Inter-Scalene Block for Acute Pain Control Following Shoulder Surgery? A Systematic Review and Meta-Analysis. *Anesthesia and Analgesia*. 2020May;130(5):1304-19. doi:10.1213/ANE.0000000000004692.
- [7] Liu S, Zhao P, Cui Y, Lu C, Ji M, Liu W et al. Effect of 5- μ g Dose of Dexmedetomidine in Combination with Intrathecal Bupivacaine on Spinal Anesthesia: A Systematic Review and Meta-Analysis. *Clinical Therapeutics*. 2020Apr;42(4):676-90. doi:10.1016/j.clinthera.2020.02.009.
- [8] Alansary AM, Badawy A, Elbeialy M. Dexmedetomidine Added to Bupivacaine Versus Bupivacaine in Transincisional Ultrasound-Guided Quadratus Lumborum Block in Open Renal Surgeries: A Randomized Trial. *Pain Physician*. 2020;23(3):271. doi:10.36076/ppj.2020/23/271.
- [9] Van Holsbeeck M, Soliman S, Van Kerkhove F, Craig J. Advanced Musculoskeletal Ultrasound Techniques: What Are the Applications? *American Journal of Roentgenology*. 2021Feb;216(2):436-45. doi:10.2214/AJR.20.22840.
- [10] Da Silva T, Mueck D, Knop C, Merkle T. Ultrasound-Guided Localization of the Radial Nerve Along the Humerus: Providing Reference Points for Safer Upper Arm Surgery. *Musculoskeletal Surgery*. 2024Jul;1-7. doi:10.1007/s12306-024-00841-1.
- [11] Gadsden JC. The Role of Peripheral Nerve Stimulation in the Era of Ultrasound-Guided Regional Anesthesia. *Anesthesia*. 2021Jan;76:65-73. doi:10.1111/anae.15257.
- [12] Althubaiti A. Sample Size Determination: A Practical Guide for Health Researchers. *Journal of General and Family Medicine*. 2023Mar;24(2):72-8. doi:10.1002/jgf2.600.
- [13] Zhao ZF, Du L, Wang DX. Effects of Dexmedetomidine as A Perineural Adjuvant for Femoral Nerve Block: A Systematic Review and Meta-Analysis. *PLOS ONE*. 2020Oct;15(10):e0240561. doi:10.1371/journal.pone.0240561.
- [14] Nazir N and Jain S. A Randomized Controlled Trial Study on the Effect of Adding Dexmedetomidine to Bupivacaine in Supraclavicular Block Using Ultrasound Guidance. *Ethiopian Journal of Health Sciences*. 2016; 26(6):561-6. doi:10.4314/ejhs.v26i6.9.
- [15] Begum MR and Hossain MA. Validity and Reliability of Visual Analogue Scale (VAS) for Pain Measurement. *Journal of Medical Case Reports and Reviews*. 2019; 2(11).
- [16] Kumar S, Choudhury B, Varikasuvu SR, Singh H, Kumar S, Lahon J et al. A Systematic Review and Meta-Analysis of Efficacy and Safety of Dexmedetomidine Combined with Intrathecal Bupivacaine Compared to Placebo. *Cureus*. 2022Dec;14(12). doi:10.7759/cureus.32425.
- [17] Sane S, Mahoori A, Tabrizi A, Koohsoltani H. Comparison of the Effect of Adding Dexmedetomidine as an Adjuvant to Bupivacaine for Postoperative Pain Management in Patients Undergoing Shoulder Rotator Cuff Repair-A Randomized Clinical Trial. *Advanced*

- Biomedical Research.2024Dec;13(1):124.doi:10.4103/abr.abr_585_24/
- [18] Ammar AS and Mahmoud KM. Ultrasound-Guided Single Injection Infraclavicular Brachial Plexus Block Using Bupivacaine Alone or Combined with Dexmedetomidine for Pain Control in Upper Limb Surgery: A Prospective Randomized Controlled Trial. Saudi Journal of Anesthesia.2012Apr;6(2):109-14.doi: 10.4103/1658-354X.97021.
- [19] Hussain A, Majeed K, Akhter A, Qureshi S, Ahsan A. Efficacy of Dexmedetomidine Combined with Bupivacaine in Supraclavicular Block Versus Bupivacaine Alone in Upper Limb Surgery.Pakistan Armed Forces Medical Journal.2023;73(3):763-6.doi: 10.51253/pafmj.v73i3.8375.
- [20] Alarasan AK, Agrawal J, Choudhary B, Melhotra A, Uike S, Mukherji A. Effect of Dexamethasone in Low Volume Supraclavicular Brachial Plexus Block: A Double-Blinded Randomized Clinical Study.Journal of Anesthesiology Clinical Pharmacology.2016 Apr; 32(2): 234-9. doi: 10.4103/0970-9185.182108.
- [21] Sane S, Shokouhi S, Golabi P, Rezaeian M, Kazemi Haki B. The Effect of Dexmedetomidine in Combination with Bupivacaine on Sensory and Motor Block Time and Pain Score in Supraclavicular Block.Pain Research and Management.2021;2021(1):8858312.doi: 10.1155/2021/8858312.