



## Original Article

## Clinical Risk Factors of Post-Surgery Hemorrhage in Patients Undergoing Tonsillectomy

Sanallah Bhatti<sup>1\*</sup>, Javed Mirdad Tarar<sup>2</sup>, Muhammad Asif Ashraf<sup>1</sup> and Mahreen Tahir<sup>1</sup>

<sup>1</sup>Department of ENT, Bakhtawar Amin Hospital, Multan, Pakistan

<sup>2</sup>Department of Thoracic Surgery, Bakhtawar Amin Trust Teaching Hospital, Multan, Pakistan

## ARTICLE INFO

**Keywords:**

Hemorrhage, Post-Tonsillectomy Hemorrhage, Tonsillectomy, Risk Factors

**How to Cite:**

Bhatti, S., Tarar, J. M., Ashraf, M. A., & Tahir, M. (2025). Clinical Risk Factors of Post-Surgery Hemorrhage in Patients Undergoing Tonsillectomy: Post-Surgery Hemorrhage in Patients Undergoing Tonsillectomy. *Pakistan Journal of Health Sciences*, 6(4), 165-169. <https://doi.org/10.54393/pjhs.v6i4.2868>

**\*Corresponding Author:**

Sanallah Bhatti  
 Department of ENT, Bakhtawar Amin Hospital,  
 Multan, Pakistan  
[sanallahdr74@gmail.com](mailto:sanallahdr74@gmail.com)

Received date: 12<sup>th</sup> February, 2025

Revised date: 25<sup>th</sup> April, 2025

Acceptance date: 27<sup>th</sup> April, 2025

Published date: 30<sup>th</sup> April, 2025

## ABSTRACT

Post-tonsillectomy hemorrhage is rare but life-threatening and poses a critical emergency. However, limited research has been conducted on adults undergoing tonsillectomy. **Objective:** To analyze the predictors of post-tonsillectomy hemorrhage in adults undergoing tonsillectomy. **Methods:** A retrospective study was conducted in the ENT and Surgery Department of Bakhtawar Amin Hospital from April 2024 to Jan 2025. A total of 150 adult patients underwent bilateral tonsillectomies for recurrent tonsillitis, obstructive sleep apnea, tonsillar hypertrophy, palmoplantar pustulosis, or IgA nephropathy. The surgery was performed according to the usual procedure under anesthesia. Age, gender, duration of surgery, obesity, antibiotic administration, smoking status, and type of analgesia were recorded as probable predictors of hemorrhage. Smoking status included current smokers, non-smokers, and former smokers who had stopped smoking at least 1 month before the procedure. **Results:** There was a significant difference between smokers and non-smokers (OR=3.52, 95% CI: 1.76-6.68,  $p<0.001$ ) and smokers and former smokers (OR=3.55, 95% CI: 1.63-7.61,  $p<0.003$ ). Male gender (OR: 4.03, 95% CI: 1.63-9.89,  $p=0.005$ ) and NSAID (OR: 7.87, 95% CI: 1.007-63.53,  $p=0.0502$ ) were significantly associated with post-operative hemorrhage. Smoking status ( $p=0.052$ ) and older age ( $p=0.005$ ) were significant risk factors in the primary hemorrhage group, and smoking status ( $p<0.001$ ) and male gender ( $p=0.010$ ) were significant in the secondary hemorrhage group. **Conclusions:** Post-tonsillectomy hemorrhage had a significant association with male gender, smoking status, and administration of NSAIDs during surgery. Therefore, it is advisable for surgeons to strongly suggest that patients maintain a healthy lifestyle before surgery and opt for other analgesics to prevent the risk of hemorrhage.

## INTRODUCTION

Tonsillectomy is a frequent procedure involving the removal of the tonsils due to tonsillitis, infections, sleep-disordered breathing, or sleep apnea in children and adults. Although the data on the Pakistani population is unavailable, the annual incidence rate in Norway is 0.16%, 0.13% in Sweden, and 0.08% in Denmark [1]. In children, it is often performed with adenoidectomy to resolve breathing and swallowing issues. Dehydration, pain, hemorrhage, nausea & vomiting, and infections are common complications after tonsillectomy [2]. Among these, post-tonsillectomy hemorrhage is rare but life-threatening and poses a critical emergency. It can cause airway obstruction and hypovolemic shock which may require surgery for management. Since the majority of tonsillectomies are

performed in children, most literature focuses on outcomes and complications in the pediatric population [3]. In children, the cause of surgery has been reported as the main cause of post-tonsillectomy hemorrhage in reviews and meta-analyses [4]. Children undergoing surgery for recurrent infections had a high rate of hemorrhage as compared to children with obstructive sleep-disordered breathing. However, hemorrhage rates did not differ with the change in surgical techniques, including coblation, cold dissection, or electrocautery [5]. An increase in age significantly enhanced the risk of hemorrhage even in pediatric samples. Only limited research has been conducted on adults undergoing tonsillectomy. The reports available are conducted on

pediatric and adult populations, which conclude that there is a high risk of hemorrhage in older age [6]. However, these results did not identify the risk factors of hemorrhage in adults specifically. We establish a hypothesis that lifestyle parameters, including surgical cause, smoking status, obesity, etc., are predictors of post-tonsillectomy hemorrhage in adults. We aim to investigate the impact of smoking status, role of surgical technique, medication and demographics on hemorrhage rates.

Post-tonsillectomy hemorrhage is a rare but serious complication that can lead to life-threatening outcomes, particularly in adult patients, where limited evidence is available compared to pediatric populations. Although several studies have identified general risk factors such as age and surgical technique, there is insufficient focused research on modifiable and demographic predictors like smoking status, NSAID use, and gender, specifically in adults. This study aimed to evaluate clinical and lifestyle-related risk factors associated with post-tonsillectomy hemorrhage in adults, with particular emphasis on identifying predictors of primary and secondary bleeding to improve perioperative risk stratification and patient safety.

## METHODS

A retrospective study was conducted in the ENT and Surgery Department of Bakhtawar Amin Trust Hospital from April 2024 to January 2025. A total of 150 adult patients undergoing bilateral tonsillectomies for recurrent tonsillitis, obstructive sleep apnea, tonsillar hypertrophy, palmpoplantar pustulosis, or IgA nephropathy were included by consecutive sampling. Adenoidectomy and/or uvulopalatopharyngoplasty were also performed in patients with sleep apnea and hypertrophy. The sample size was calculated by keeping a 50% population proportion, 95% confidence interval, 80% power and precision of  $\pm 5\%$  among 245 population size. Patients who underwent tonsillectomy for tumor or quinsy, or those undergoing head and neck surgery for other conditions or hematologic disease, were excluded. All patients provided their informed consent to become a part of the study. The ethical board of the hospital approved the study Ref No.3294/BAMTH. The surgery was performed according to the usual procedure under anesthesia. After intubating the patient, Crowe-Davis mouth gag to secure the tonsil site. A mucosal incision was made, and the tonsils were released from their capsule using a dissector. Bipolar or mono-polar electrocautery was used to maintain hemostasis. No other equipment was used. Where electrocautery was ineffective in controlling bleeding, absorbable sutures were employed to ligate bleeding points. IV antibiotics were given intraoperatively and postoperatively. Fentanyl and/or remifentanyl were also administered during surgery, and acetaminophen after surgery as analgesics. Post-tonsillectomy hemorrhage was categorized based on

Windfuhr's classification. In addition, hemorrhage within 24 hours after the surgery was called primary hemorrhage, and hemorrhage after 24–48 hours was called secondary hemorrhage. Age, gender, duration of surgery, obesity, antibiotic administration, smoking status, and type of analgesia were recorded as probable predictors of hemorrhage. Smoking status included current smokers, non-smokers, and former smokers who had stopped smoking at least 1 month before the procedure. All data were analyzed by SPSS version 24.0. Descriptive analysis was performed on quantitative variables and was presented by frequency and percentage. Univariate analysis was performed by the Mann-Whitney U test and multivariate analysis was performed by Fisher's exact test to recognize risk factors of post-operative bleeding. A p-value less than 0.05 was considered significant.

## RESULTS

A total of 150 adult patients undergoing tonsillectomy were included for analysis. Among the study subjects, 114 patients (76%) were male and 36 (24%) were female. The average age of patients was  $32.2 \pm 9.8$  years. 33 patients (22%) were smokers while 72 patients (48%) were non-smokers. 111 (74%) patients underwent surgery for recurrent tonsillitis, 21 (14%) for IgA nephropathy, 1 (0.7%) for palmpoplantar pustulosis, 15 (10%) for obstructive sleep apnea, and 3 (2%) for tonsillar hypertrophy. 9 patients (6%) underwent adenoidectomy and 15 patients (10%) underwent uvulopalatopharyngoplasty. 33 patients (22%) among 150 patients experienced post-tonsillectomy hemorrhage with 28 (85%) being grade 1, 2 (6%) being grade 2, 3 (9%) being grade 3, and 117 (78%) patients had no bleeding. Grade 2 and 3 patients required intervention to maintain hemostasis, and grade 3 patients also underwent additional surgeries. 27 (24.4%), 4 patients (26.7%), and 2 (14.3%) patients had a hemorrhage in patients undergoing tonsillectomy for recurrent tonsillitis, sleep apnea, and IgA nephropathy, respectively ( $p=0.277$ ) (Table 1).

**Table 1:** Surgical Indication of Tonsillectomies and Classification of Post-Tonsillectomy Hemorrhage (n=150)

Surgical Indication	n (%)	Post-Tonsillectomy Hemorrhage	Primary Hemorrhage	Secondary Hemorrhage
Recurrent Tonsillitis	111 (74%)	27 (24.4%)	4 (3.6%)	23 (20.8%)
IgA Nephropathy	21 (14%)	2 (14.3%)	-	2 (14.3%)
Palmpoplantar Pustulosis	1 (0.7%)	-	-	-
Sleep Apnea	15 (10%)	4 (26.7%)	1 (6.7%)	3 (20%)
Tonsillar Hypertrophy	3 (2%)	-	-	-

The study analyzed the risk factors of postoperative hemorrhage by univariate analysis. The smoking status differed significantly between total cases ( $p<0.001$ ),

primary hemorrhage group ( $p=0.05$ ), and secondary hemorrhage group ( $p<0.001$ ) as compared to the no bleeding group. Male gender was more prevalent in the total hemorrhage group ( $p<0.001$ ) and secondary hemorrhage group ( $p=0.005$ ) as compared to the no bleeding group. The primary hemorrhage group had a significantly higher average age ( $p=0.005$ ) (Table 2).

**Table 2:** Univariate Analysis

Risk Factors	No Post-Tonsillectomy Hemorrhage (n=117)	Post-Tonsillectomy Hemorrhage (n=33)	p-value	Primary Hemorrhage (n=5)	p-value	Secondary Hemorrhage (n=28)	p-Value
Average age	32.3	33.5	0.512	44.1	0.005**	32.2	0.915
<b>Gender</b>							
Male	84 (71.8%)	30 (91%)	<0.001***	5 (100%)	0.093	25 (89.5%)	0.005**
Female	33 (28.2%)	3 (9%)		-		3 (10.5%)	
<b>BMI</b>							
25 Or More	39 (33.4%)	30 (91%)	1.000	3 (60%)	0.524	8 (28.6%)	0.716
Less Than 25	78 (66.7%)	3 (9%)		2 (40%)		20 (71.4%)	
Current Smokers	19 (16.3%)	14 (42.5%)	<0.001***	3 (60%)	0.050*	11 (39.3%)	<0.001***
Duration of Surgery (Minutes)	79.1	78.0	1.000	64.1	0.202	79.3	0.571
<b>Postoperative Antibiotics</b>							
Oral intake	87 (74.4%)	23 (69.9%)	0.612	4 (80%)	1.000	19 (68%)	0.533
Injection	30 (25.6%)	10 (30.1%)		1 (20%)		9 (32%)	
<b>Analgesia</b>							
Non-Steroidal Anti-Inflammatory Drugs	106 (90.6%)	32 (97%)	0.055	5 (100%)	1.000	27 (96.4%)	0.096
Acetaminophen	11 (9.4%)	1 (3%)		-		1 (3.6%)	

\* $p<0.050$ , \*\* $p<0.010$  and \*\*\* $p<0.001$

Results show a multivariate analysis of post-tonsillectomy hemorrhage. There was a significant difference between smokers and non-smokers ( $OR=3.52$ , 95% CI: 1.76–6.68,  $p<0.001$ ) and smokers and former smokers ( $OR=3.55$ , 95% CI: 1.63–7.61,  $p<0.003$ ). Male gender ( $OR: 4.03$ , 95% CI: 1.63–9.89,  $p=0.005$ ) and NSAID ( $OR: 7.87$ , 95% CI: 1.007–63.53,  $p=0.050$ ) were significantly associated with post-operative hemorrhage. Smoking status ( $p=0.052$ ) and older age ( $p=0.005$ ) were significant risk factors in the primary hemorrhage group, and smoking status ( $p<0.001$ ) and male gender ( $p=0.010$ ) were significant in the secondary hemorrhage group (Table 3).

**Table 3:** Multivariate Analysis

Risk Factors	Post-Tonsillectomy Hemorrhage		Primary Hemorrhage		Secondary Hemorrhage	
	OR (95% CI)	p-value	OR (95% CI)	p-value	OR (95% CI)	p-value
Age	0.99 (1.0-1.08)	0.191	1.09 (1.05-1.30)	0.005*	0.99 (1.0-1.06)	0.833
Male Gender	4.03 (1.63-9.89)	0.005**	-	-	3.47 (1.42-9.03)	0.010**
Obesity	0.59 (0.30-1.18)	0.225	0.61 (0.11-3.32)	0.492	0.59 (0.28-1.17)	0.223
Current Smokers vs Non-Smokers	3.52 (1.76-6.68)	<0.001**	7.18 (1.05-49.88)	0.052*	3.29 (1.70-6.64)	<0.001**
Duration of Surgery	0.98 (1.01-1.03)	0.678	1.01 (0.95-1.05)	0.213	0.99 (1.0-1.02)	0.957
Antibiotics	0.75 (0.49-1.60)	0.615	1.01 (0.20-5.67)	1.021	0.77 (0.39-1.62)	0.600
NSAIDS	7.87 (1.007-63.53)	0.050*	-	-	7.02 (0.94-54.44)	0.100

\* $p<0.050$ , \*\* $p<0.010$  and \*\*\* $p<0.001$

## DISCUSSION

This study was conducted to analyze the predictors of postoperative hemorrhage in adults undergoing tonsillectomy. The results revealed that male gender, smoking status, and administration of NSAIDs during surgery were independently associated with postoperative bleeding. The overall risk of hemorrhage and primary and secondary bleeding was increased in current smokers. Other studies have also reported smoking as a dependent predictor of hemorrhage in tonsillectomy patients [7–9].

However, there is no data regarding the association of hemorrhage with former smokers. In present study, it was showed that a history of smoking by former smokers was not related to an increased risk of hemorrhage in comparison to non-smokers. However, taking into account the duration of cessation in former smokers can help yield better results. Since inflammation and infection are side effects of smoking, post-operative wound healing is also worse in smokers, which increases the likelihood of

hemorrhage [10]. Additionally, these patients also have increased sputum and mucus production, which can stimulate coughing postoperatively and bleeding eventually [11]. Hence, patients must advise abstinence or cessation of smoking before the procedure to avoid critical emergencies. Gender is a significant independent predictor of overall hemorrhage and secondary bleeding, with increased incidence in males. Previous studies investigating the link between gender and hemorrhage risk have also reported the same findings, which can be explained by the fact that estrogen contributes to faster healing and prevents inflammation, therefore, women experience fewer complications [7, 12]. However, gender has not been significant in children undergoing tonsillectomy because sex hormones are not differentiated before puberty. Smoking is also more prevalent in men than in women in Pakistan, hence increasing the risk of hemorrhage in men, however, smoking status and male gender were independent predictors in present study. Older age was also an independent predictor of primary hemorrhage since tonsillar inflammation can last for a long time in older patients. As age increases, blood vessels weaken, increasing the likelihood of bleeding shortly after the procedure. Overall, hemorrhage risk was also significantly linked to intraoperative administration of nonsteroidal anti-inflammatory drugs (NSAIDs). Literature found that NSAIDs increased the risk of bleeding after tonsillectomy as these medications block cyclooxygenase, leaving an antiplatelet effect [13]. McLean *et al.*, concluded that patients administered NSAIDs were more likely to require surgery to treat post-tonsillectomy hemorrhage, but So *et al.*, drew contrasting results that there was an association between post-operative bleeding and its surgical treatment and use of NSAIDs [14, 15]. Although there is no definite view about NSAIDs being a risk factor for hemorrhage, the present study supports the results of McLean *et al.*, [14]. Other probable risk factors, including duration of surgery, obesity, and antibiotic use, were not related to the incidence of hemorrhage. 74% of patients in present study underwent surgery for recurrent tonsillitis. Research shows that postoperative hemorrhage in patients who underwent surgery for recurrent tonsillitis is linked to prolonged inflammation [16, 17]. However, there was no significant difference in surgical indications in current study. Since we included adult patients, tonsillar inflammation may have occurred with other risk factors like smoking, but may have been asymptomatic due to which we could not consider surgical indication as a predictor of hemorrhage. The incidence of post-tonsillectomy hemorrhage was 22% in current study, which is higher than the rate reported by previous studies, i.e. 2.2%-10% [18, 19]. 9% of patients in current study required surgeries to

resolve the hemorrhage, which is also higher than 1-6% reported in previous studies [20].

This study is limited by its retrospective, single-center design and relatively small sample size, which may affect the generalizability of the findings. In addition, potential confounding factors such as surgical technique variation and unmeasured patient comorbidities may influence results. Future research should involve large-scale, multicenter prospective studies to validate these risk factors more robustly. It is also recommended to develop standardized perioperative guidelines focusing on smoking cessation and NSAID alternatives to reduce the risk of post-tonsillectomy hemorrhage in high-risk patients.

## CONCLUSIONS

It was concluded that post-tonsillectomy hemorrhage had a significant association with male gender, smoking status, and administration of NSAIDs during surgery. Therefore, it is advisable for surgeons to strongly suggest that patients maintain a healthy lifestyle before surgery and opt for other analgesics to prevent the risk of hemorrhage.

## Authors' Contribution

Conceptualization: SB, JMT

Methodology: MAA, MT

Formal analysis: JMT

Writing and Drafting: MT

Review and Editing: MAA, MT, SB, JMT

All authors approved the final manuscript and take responsibility for the integrity of the work

## Conflicts of Interest

All the authors declare no conflict of interest.

## Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

## REFERENCES

- [1] Stalfors J, Ovesen T, Bertelsen JB, Bugten V, Wennberg S, Sunnergren O. Comparison of Clinical Practice of Tonsil Surgery from Quality Register Data from Sweden and Norway and One Clinic in Denmark. *BioMed Journal Open*. 2022 Apr;12(4):e056551. doi:10.1136/bmjopen-2021-056551.
- [2] Patel SD, Daher GS, Engle L, Zhu J, Slonimsky G. Adult Tonsillectomy: An Evaluation of Indications and Complications. *American Journal of Otolaryngology*. 2022 May;43(3):103403. doi:10.1016/j.amjoto.2022.103403.
- [3] Dhaduk N, Rodgers A, Govindan A, Kalyoussef E. Post-Tonsillectomy Bleeding: A National Perspective. *Annals of Otolaryngology, Rhinology and Laryngology*. 2021 Aug;130(8):941-7. doi:10.1177/0003489420987438.

- [4] Gutierrez JA, Shannon CM, Nguyen SA, Labadie RF, White DR. The Impact of Surgical Indication On Posttonsillectomy Hemorrhage: A Systematic Review and Meta-Analysis. *Otolaryngology-Head and Neck Surgery*.2023Oct;169(4):780-91.doi:10.1002/ohn.339.
- [5] Çelikal Ö, Eroğlu E, Günaydın RÖ. Post-Tonsillectomy Hemorrhage in Pediatric Patients: Comparison of Age Groups and Surgical Techniques. *Allergy*.2023Aug;6(2):45-8.doi:10.5152/ejra.2023.23091.
- [6] Öcal B, Günay MM, Keseroğlu K, Mutlu M, Akyıldız İ, Saka C et al. Risk Factors of Post-Tonsillectomy Bleeding and Differences Between Children and Adults: Implications for Risk Assessment. *Turkish Archives of Otorhinolaryngology*.2025Jan;62(3):81.doi:10.4274/tao.2024.2023-10-2.
- [7] Fushimi K, Gyo K, Okunaka M, Watanabe M, Sugihara A, Tsuzuki K. Analysis of Risk Factors for Post-Tonsillectomy Hemorrhage in Adults. *Auris Nasus Larynx*.2023Jun;50(3):389-94.doi:10.1016/j.anl.2022.10.002.
- [8] Besiashvili N, Datikashvili-David IG, Gakharia T. Evaluation and Risk Factor Analysis of Post-tonsillectomy Hemorrhage in an Adult Population: An Experience From a National Ear, Nose, and Throat (ENT) Center in Georgia. *Cureus*.2024Sep;16(9).doi:10.7759/cureus.68371.
- [9] Krkalic A, Mandra H, Joguncic A. Tonsillectomy and the Risk of Post-Tonsillectomy Hemorrhage: A Retrospective Cross-Sectional Study. *Sarajevo Medical Journal*.2024Aug;1(1):5.doi:10.70119/000224.
- [10] KC S, Dongol SR, Bhattarai S, Mahaseth RK. Long Term Results of Adult Tonsillectomy for Recurrent Tonsillitis at a Tertiary Referral Hospital in Nepal. *Journal of Institute of Medicine Nepal*.2021Aug;43(2).doi:10.59779/jiomnepal.1170.
- [11] Arif M, Noor A, Anwar K, Khan S. Frequency of Post-Tonsillectomy Secondary Hemorrhage. *Khyber Journal of Medical Sciences*.2024Oct;17(4):223.doi:10.70520/kjms.v17i4.241.
- [12] El-magd EA, El-Tahan AR, Mohammed SA, Khalifa ME. Risk Factors for Secondary Post Tonsillectomy Hemorrhage. *International Journal of Health Sciences*.2022;6(S4):10976-86.doi:10.53730/ijhs.v6nS4.11168.
- [13] Jacobson A, Mack D, Herrera G, Bowe SN, Highland KB, Patzkowski MS. Incidence of Surgically Managed Post-Tonsillectomy Hemorrhage Associated with NSAID Prescribing for Postoperative Pain Management. *Military Medicine*.2024Sep;189(9-10):e1955-9.doi:10.1093/milmed/usae194.
- [14] McLean JE, Hill CJ, Riddick JB, Folsom CR. Investigation of Adult Post-Tonsillectomy Hemorrhage Rates and the Impact of NSAID Use. *The Laryngoscope*.2022May;132(5):949-53.doi:10.1002/lary.29844.
- [15] So V, Radhakrishnan D, MacCormick J, Webster RJ, Tsampalieros A, Zitikyte G et al. Does Celecoxib Prescription for Pain Management Affect Post-Tonsillectomy Hemorrhage Requiring Surgery? A Retrospective Observational Cohort Study. *Anesthesiology*.2024Jul;141(2):313-25.doi:10.1097/ALN.0000000000005032.
- [16] Gross JH, Lindburg M, Kallogjeri D, Molter M, Molter D, Lieu JE. Predictors of Occurrence and Timing of Post-Tonsillectomy Hemorrhage: A Case-Control Study. *Annals of Otolaryngology, Rhinology and Laryngology*.2021 Jul;130(7):825-32.doi:10.1177/0003489420978010.
- [17] Slouka D, Čejková Š, Hanáková J, Hrabačka P, Kormunda S, Kalfeřt D et al. Risk of Postoperative Bleeding in Tonsillectomy for Peritonsillar Abscess, as Opposed to in Recurrent and Chronic Tonsillitis—A Retrospective Study. *International Journal of Environmental Research and Public Health*.2021 Feb;18(4):1946. doi:10.3390/ijerph18041946.
- [18] Schafer A, Worobetz N, Lukens J, Bourgeois T, Onwuka A, Elmaraghy C et al. Assessing the Relationship Between Infection Frequency and Risk of Post-Tonsillectomy Hemorrhage. *Annals of Otolaryngology, Rhinology and Laryngology*. 2023 Nov; 132(11): 1424-9. doi:10.1177/00034894231159328.
- [19] Faiq TN and Ghareeb OA. A Review on the Actual Indications for Tonsillectomy in Adults and Potential Postoperative Complications. *World Bulletin of Public Health*. 2022 Jan; 6: 46-50.
- [20] Mohammadpour-Maleki A and Rasouljan B. Post-Tonsillectomy Hemorrhage: A Seven-Year Retrospective Study. *Iranian Journal of Otorhinolaryngology*.2021 Sep; 33(118): 311. doi:10.22038/ijorl.2021.54962.2882.