



## Original Article



# Septoplasty in Focus: Clinical Outcomes and Complication Patterns in A Pakistani Cohort

Naheed Yaqoob<sup>1</sup>, Sajjad Yousuf<sup>1\*</sup>, Akhtar Ali Lakhari<sup>1</sup>, Arslan Ali<sup>1,2</sup>, Amrat Kumar<sup>1,2</sup> and Faizan Khan<sup>1</sup>

<sup>1</sup>Department of ENT, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan

<sup>2</sup>Department of ENT, ISRA University, Hyderabad, Pakistan

## ARTICLE INFO

**Keywords:**

Septal Deviation, Septoplasty, Nasal Obstruction, Complications, Hemorrhage, Pakistan, ENT Surgery

**How to Cite:**

Yaqoob, N., Yousaf, S., Lakhari, A. A., Ali, A., Kumar, A., & Khan, F. (2025). Septoplasty in Focus: Clinical Outcomes and Complication Patterns in A Pakistani Cohort: Septoplasty in Focus: Clinical Outcomes and Complication Patterns. *Pakistan Journal of Health Sciences*, 6(7), 271-275. <https://doi.org/10.54393/pjhs.v6i7.3343>

**\*Corresponding Author:**

Sajjad Yousuf  
 Department of ENT, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan  
[sajjadzardari55@gmail.com](mailto:sajjadzardari55@gmail.com)

Received Date: 2<sup>nd</sup> June, 2025

Revised Date: 21<sup>st</sup> July, 2025

Acceptance Date: 26<sup>th</sup> July, 2025

Published Date: 31<sup>st</sup> July, 2025

## ABSTRACT

Septal deviation is a common cause of nasal obstruction, and septoplasty is the preferred corrective procedure. Despite its widespread use, septoplasty is associated with several intraoperative and postoperative complications, which remain underreported in the Pakistani population. **Objectives:** To determine the frequency and nature of complications following septal surgery at a tertiary care setting. **Methods:** An observational cohort study was conducted at the Department of Otorhinolaryngology, Civil Hospital Hyderabad and LUMHS Jamshoro from March 2020 to February 2021. A total of 174 patients undergoing septal surgery were enrolled using non-probability consecutive sampling. Data on demographics, presenting complaints, surgical procedures, and postoperative complications were collected using structured questionnaires. All patients were followed monthly for six months postoperatively. Data were analyzed using SPSS version 21.0. **Results:** The mean age of participants was  $27.5 \pm 5.2$  years, with 62.1% being male. The most common intraoperative complication was hemorrhage, observed in 49 patients (28.2%). Postoperative complications included septal hematoma in 18 patients (10.3%), infection in 17 patients (9.8%), and septal perforation in 12 patients (6.9%). A statistically significant association was observed between resection of the deviated septum and both immediate and delayed complications ( $p=0.035$ ), indicating that this procedure carried a higher risk compared to others. **Conclusions:** Septoplasty, particularly when involving resection of the deviated septum, was associated with a higher rate of both immediate and delayed complications. These findings highlight the need for careful surgical technique selection and structured postoperative monitoring to minimize morbidity and improve patient outcomes in clinical ENT practice.

## INTRODUCTION

Nasal septal deviation (NSD) represents a displacement of the nasal septum affecting approximately 80% of the global population, although symptomatic cases are fewer [1]. This anatomical deformity often leads to chronic nasal obstruction, compromised ventilation, and reduced quality of life [2, 3]. When medical management such as nasal steroids or decongestants, fails, Septoplasty becomes the treatment of choice [4]. Despite its frequent application, septoplasty is associated with known complications, including hemorrhage, hematoma, perforation, infection, and adhesions [5, 6]. Recent refinements such as endoscopic techniques and quilting sutures have been

introduced to reduce these risks and enhance recovery [7, 8]. However, in Pakistan, systematic data on complication patterns following septoplasty remain limited. A few single-center studies have explored surgical approaches and postoperative outcomes, such as a comparative study from Lahore showing shorter operative times for endonasal septoplasty compared to external approaches [9], and the use of nasal splints significantly reducing adhesions [10-14]. Another evaluation from Jinnah Hospital endorsed endoscopic techniques for enhanced safety [11]. Yet, these findings remain fragmented, and standardized outcome monitoring across institutions is



lacking. Given this context, there is a clear need to generate robust, locally relevant data on the complications of septal surgery.

Nasal septal deviation is a common anatomical condition that frequently leads to nasal obstruction and reduced quality of life, with septoplasty being the standard surgical treatment. Although the procedure is widely performed, it is still associated with various intraoperative and postoperative complications that may affect surgical outcomes. While international studies have documented these complication patterns, limited systematic evidence is available from Pakistani clinical settings regarding the frequency and types of complications following septoplasty. This lack of comprehensive local data highlights the need for further investigation to guide safer surgical practices and improve postoperative patient management. This study aims to determine the frequency and types of complications associated with septoplasty at a tertiary care center in Hyderabad, with the goal of informing surgical practice and optimizing patient care.

## METHODS

This observational cohort study was conducted in the Department of Otorhinolaryngology (ENT) at Liaquat University Hospital, Hyderabad, from March 2020 to February 2021. The hospital was a tertiary care teaching facility affiliated with Liaquat University of Medical and Health Sciences (LUMHS), Jamshoro. A total of 174 patients undergoing septal surgery were included. The sample size was calculated using the OpenEpi sample size calculator, based on a hypothesized complication rate of 13% [6], with a 5% margin of error and a 95% confidence level. The study was approved by the Ethical Review Committee of LUMHS, Jamshoro (Approval Letter No. LUMHS/REC/-791). Informed written consent was obtained from all participants in Urdu or a local regional language, ensuring full comprehension of the study's nature, risks, and benefits. Patient confidentiality was maintained throughout the research process. Participants were enrolled through a non-probability consecutive sampling technique. The inclusion criteria comprised all patients of any age and gender who presented with nasal airway obstruction due to septal deviation and were scheduled for corrective septal surgery (septoplasty). Patients were excluded if they had a history of previous septal surgery, external nasal deformities, associated nasal pathologies such as nasal polyps or rhinoliths, or known hemostatic disorders. Data on demographic and clinical variables were collected using a structured, pre-tested questionnaire administered at the time of patient admission. Demographic data included age and gender. Clinical data included the primary presenting complaint, physical findings, the type of surgical procedure performed, and

complications encountered. All surgical procedures including Septoplasty, septal scoring, partial inferior turbinectomy, and combined techniques were performed by ENT surgeons with experience at least of 5 years, using standardized operative protocols. Septoplasty involved Killian's incision, elevation of mucoperichondrial flaps, resection or repositioning of deviated segments, and suturing with absorbable material. Scoring was performed in mild deviations without cartilage removal. Partial inferior turbinectomy was done using turbinate scissors or micro-debriders to reduce turbinate size. All patients received preoperative antibiotics, and hemostasis was achieved using cautery or Gelfoam. Postoperative care followed a uniform regimen, and postoperative complications were documented using a standardized follow-up proforma. Patients were followed for six months postoperatively, with clinical evaluations scheduled monthly at the ENT outpatient department. Assessments were conducted monthly up to 6 months after surgery, during which each patient underwent anterior rhinoscopy and nasal endoscopy to identify potential complications. Immediate complications were defined as those occurring intraoperatively or within 48 hours' post-surgery (e.g., hemorrhage, septal perforation), while delayed complications included those identified during any of the subsequent follow-up visits (e.g., hematoma, abscess, adhesions, anosmia, infection, or structural deformities). Data analysis was performed using SPSS version 21.0. Quantitative variables such as age were expressed as mean and standard deviation (Mean  $\pm$  SD), while qualitative variables such as gender, surgical procedure type, and complications were summarized as frequencies and percentages (n, %). The Chi-square test was applied to determine the association between the type of surgical procedure and the incidence of complications. A p-value of less than 0.050 was considered statistically significant.

## RESULTS

A total of 174 patients undergoing septal surgery were included in the study. The mean age of the participants was  $27.54 \pm 5.21$  years, indicating a predominantly young adult population. There was a clear male predominance (62.1%) and a higher proportion of patients from urban areas (73%). Most patients belonged to the lower socioeconomic class, and nasal obstruction, either unilateral or bilateral, was the most common clinical presentation, followed by facial discomfort and headache (Table 1).

**Table 1:** Demographics and Clinical Profile of Patients Undergoing Septal Surgery (n=174)

Variable	Frequency (%), Mean $\pm$ SD
Mean Age (yrs)	27.54 $\pm$ 5.21
<b>Gender</b>	
Male	108 (62.1%)

Female	66 (37.9%)
<b>Residence</b>	
Urban	127 (73%)
Rural	47 (27%)
<b>Presenting Complaint</b>	
Facial Discomfort	18 (10.3%)
Unilateral Nasal Obstruction	51 (29.3%)
Bilateral Nasal Obstruction	66 (37.9%)
Frontal Headache	19 (10.9%)
Nose Bleeds	11 (6.3%)
Snoring	9 (5.2%)

With regard to postoperative outcomes, complications

were stratified by type and time of onset. The most common intraoperative complication was hemorrhage, reported in 28.2% of patients. Among early postoperative complications, septal hematoma was most frequent in the first month (10.3%), with rates decreasing in subsequent months. Septal perforation was reported in 6.9% intraoperatively, with a small number of new cases (1.1%) emerging within the first postoperative month. Structural deformities, anosmia/hyposmia, and infections were distributed across the follow-up period, while adhesions and septal abscesses tended to occur later, though at lower frequencies overall (Table 2).

**Table 2:** Incidence of Complications by Time of Onset (n=174)

Complications	During Surgery	Month 1	Month 2	Month 3	Month 4	Month 5	Month 6
Hemorrhage	49 (28.2%)	-	-	-	-	-	-
Septal Hematoma	-	18 (10.3%)	11 (6.3%)	4 (2.3%)	1 (0.6%)	-	-
Adhesion	-	-	-	-	1 (0.6%)	3 (1.7%)	5 (2.9%)
Septal Perforation	12 (6.9%)	2 (1.1%)	-	-	-	-	-
Structural Deformity	-	9 (5.2%)	2 (1.1%)	-	-	-	-
Anosmia / Hyposmia	-	2 (1.1%)	1 (0.6%)	1 (0.6%)	1 (0.6%)	-	1 (0.6%)
Infection	-	17 (9.8%)	12 (6.9%)	9 (5.2%)	-	-	-
Septal Abscess	-	1 (0.6%)	1 (0.6%)	2 (1.1%)	-	-	-

A comparative analysis of complication rates by type of surgical procedure revealed a statistically significant association between resection of the deviated septum and both immediate (9.8%) and delayed complications (16.7%) (p=0.035). In contrast, scoring of the septum, combined resection and scoring, and partial inferior turbinectomy were not significantly associated with complication frequency, despite having comparatively higher percentages in some cases (p=0.410, 0.270, and 0.190 respectively). These findings suggest that resection alone may carry a higher risk of adverse outcomes compared to other techniques (Table 3).

**Table 3:** Type of Procedure Vs. Incidence of Complications (n=174)

Procedures	Immediate Complications (%)	Immediate CI (95%)	Delayed Complications (%)	Delayed CI (95%)	p-value
<b>Resection of Deviated Septum</b>					
Present	17 (9.8%)	6.2%-15.1%	29 (16.7%)	11.9%-22.9%	0.035
Absent	157 (90.2%)		145 (83.3%)		
<b>Scoring of Septum</b>					
Present	5 (2.9%)	1.2%-6.5%	9 (5.2%)	2.7%-9.5%	0.410
Absent	169 (97.1%)		165 (94.8%)		
<b>Resection &amp; Scoring of Nasal Septum</b>					
Present	18 (10.3%)	6.6%-15.8%	31 (17.8%)	12.8%-24.2%	0.270
Absent	156 (89.7%)		143 (82.2%)		
<b>Partial Inferior Turbinectomy</b>					
Present	21 (12.1%)	8.0%-17.7%	35 (20.1%)	14.8%-26.7%	0.190
Absent	153 (87.9%)		139 (79.9%)		

## DISCUSSION

This study evaluated 174 patients undergoing septal surgery and found that hemorrhage was the most frequent intraoperative complication (28.2%), while septal hematoma (10.3%), infection (9.8%), and septal perforation (6.9%) were the most commonly observed postoperative complications over six months of follow-up. Resection of the deviated septum was significantly associated with

higher complication rates (p=0.035), whereas other procedures, such as scoring or turbinectomy, did not show statistically significant associations. The mean age of the study cohort was 27.54±5.21 years, with a male predominance (62.1%). The most common presenting complaint was nasal obstruction (56.3%), followed by frontal headache (24.7%) and facial discomfort (19.0%).

These symptoms, which collectively affect patients' quality of life, have been consistently reported as the primary indications for septal surgery in both regional and international literature [15]. Analysis of septal deviation revealed varied presentations: unilateral nasal obstruction was noted in 29.3%, bilateral nasal obstruction in 37.9%, with lesser frequencies of other complaints such as facial discomfort (10.3%), frontal headache (10.9%), nosebleeds (6.3%), and snoring (5.2%). These clinical patterns support the necessity for individualized surgical approaches based on patient symptomatology and anatomical variations [6]. The incidence of complications following septal surgery was meticulously documented over a six-month follow-up period. Hemorrhage was the most common intraoperative event, comparable to previous Pakistani studies reporting similar rates between 25–30% [16]. Septal hematoma and infection were also frequent, particularly during the first month. Although complications such as adhesions, perforations, and anosmia were less prevalent, their potential impact on long-term surgical success underscores the importance of meticulous technique and postoperative care [17]. Current finding that resection of the deviated septum was significantly associated with complications aligns with studies suggesting that more invasive techniques increase tissue trauma and thus complication risk. For instance, similar associations have been reported in international literature, linking extensive resection with mucosal disruption and delayed healing [18, 19]. Moreover, current results concur with a recent European multicenter study, which emphasized that complication rates can be minimized through careful case selection and preference for minimally invasive techniques [20].

This study has certain limitations that should be considered while interpreting the findings. The research was conducted at a single tertiary care center using a non-probability sampling technique, which may limit the generalizability of the results. Additionally, the study focused mainly on short-term complications and did not evaluate long-term functional outcomes such as patient quality of life or symptom improvement. Future multicenter studies with larger sample sizes and longer follow-up periods are recommended to better assess the long-term safety and effectiveness of different septoplasty techniques and to establish standardized surgical protocols.

## CONCLUSIONS

Septoplasty is effective for nasal obstruction but carries a risk of complications, especially with resection techniques. Hemorrhage, hematoma, and infection were common. Findings support careful surgical selection and structured follow-up to improve outcomes and guide clinical practice.

## Authors' Contribution

Conceptualization: NY

Methodology: NY, SY, AAL, AA, FK

Formal analysis: AK, FK

Writing and Drafting: SY, AAL

Review and Editing: SY, AAL, NY, AK, FK

All authors approved the final manuscript and take responsibility for the integrity of the work

## Conflicts of Interest

All the authors declare no conflict of interest.

## Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

## REFERENCES

- [1] Cleveland Clinic. Deviated Septum [Internet]. Cleveland Clinic; 2023 [cited 2025 Jun 11]. Available from: <https://my.clevelandclinic.org/health/diseases/16924-deviated-septum>.
- [2] Fageeh YA, Basurrah MA, ALazwari KD, AlAmri MA, AlJuaid WM, AlHumaidi AA et al. Prevalence of Nasal Obstruction and Its Impact on Quality of Life in Saudi Arabia. *Journal of Family Medicine and Primary Care*. 2024Feb;13(2):572-8. doi:10.4103/jfmpc.jfmpc\_48\_2\_23.
- [3] Hsu DW and Suh JD. Anatomy and Physiology of Nasal Obstruction. *Otolaryngologic Clinics of North America*. 2018Oct;51(5):853-65. doi:10.1016/j.otc.2018.05.001.
- [4] Carrie S, Fouweather T, Homer T, O'Hara J, Rousseau N, Rooshenas L et al. Effectiveness of Septoplasty Compared to Medical Management in Adults with Obstruction Associated with A Deviated Nasal Septum: The NAIROS RCT. *Health Technology Assessment (Winchester, England)*. 2024Mar;28(10):1. doi: 10.3310/MVFR4028.
- [5] Chhatbar YB, Sharma S, Patel PM, Bavavva DD, Patel VB. Complications and Management of Endoscopic Septoplasty at A Tertiary Care Center in India. *International Journal of Scientific Research*. 2020 Nov; 9(11): 82-84. doi: 10.36106/ijsr/6417928.
- [6] Dąbrowska-Bień J, Skarżyński PH, Gwizdalska I, Łazęcka K, Skarżyński H. Complications in Septoplasty Based on A Large Group of 5639 Patients. *European Archives of Oto-Rhino-Laryngology*. 2018Jul;275(7):1789-94. doi:10.1007/s00405-018-4990-8.
- [7] Besharah BO, Alharbi HA, Suliman OA, Althobaiti HK, Mogharbel AM, Muathen SH. Endoscopic Septoplasty Versus Conventional Septoplasty for Nasal Septum Deviation: A Systematic Review and Meta-Analysis of

- Randomized Clinical Trials. *Annals of Medicine and Surgery*. 2023 Aug; 85(8): 4015-25. doi: 10.1097/MS9.0000000000000984.
- [8] Majeed SA and Saeed BM. The Efficacy of Septal Quilting Sutures Versus Nasal Packing in Septoplasty. *Indian Journal of Otolaryngology and Head and Neck Surgery*. 2022 Oct; 74(Suppl 2):1713-7. doi:10.1007/s12070-021-02865-6.
- [9] Awan NU, Azeem M, Cheema KM. Comparison of Endonasal and External Septoplasty for Treatment of Buckled Caudal Septal Deviation. *Journal of College of Physicians and Surgeons Pakistan*. 2021 Jan; 31(1): 51-4. doi: 10.29271/jcpsp.2021.01.51.
- [10] Asif A, Ahmed A, Rafique U, Azam K, Malik NK, Manzoor M. Role of Nasal Splints in Prevention of Nasal Adhesions following Septal Surgery at CMH Lahore: A Comparative Cross-Sectional Study. *Life and Science*. 2024 Aug; 5(3): 06-. doi: 10.37185/LnS.1.1491.
- [11] Haider H, Aziz B, Wahab U, Rehman AU, Aleem Z, Umar M et al. Analogical Evaluation of Endoscopic Septoplasty in Posterior Septal Deviation. *Pakistan Journal of Medical and Health Sciences*. 2022 Dec; 16(11): 106-. doi: 10.53350/pjmhs20221611106.
- [12] Raza MS, Malik MN, Butt NU, Haider M, Ashfaq AH, Umar M et al. Incidence of Deviated Nasal Septum in Patients Presented in Outpatient Department. *Resident Journal of Rawalpindi Medical University*. 2022 Nov; 3(11).
- [13] Dawn News. ENT Specialists Concerned Over Rise in Nasal, Throat Ailments [Internet]. Karachi: Dawn; 2024 Jun. [cited 2025 Jun 11].
- [14] Afridi AU, Ali KU, Bhatti HM, Gohar MS, Slahuddin SA, Ahmad WA. Improvement in Quality of Life After Septoplasty. *Pakistan Journal of Medical and Health Sciences*. 2022; 638-40. doi:10.53350/pjmhs221610638.
- [15] Muthubabu K, Srinivasan MK, Thejas SR, Sindu M, Vinayak R, Gayathri CS. Quality of Life in Patients with Nasal Septal Deviation After Septal Correction. *Indian Journal of Otolaryngology and Head and Neck Surgery*. 2019 Nov; 71(Suppl 3):2219-24. doi:10.1007/s12070-019-01689-9.
- [16] Touihmi S, Sadki A, Lachkar A, Elayoubi F, Serraj K, Ghailan MR. Slow Progressing Localized Tumor-Like Pharyngeal Mass: What Is Your Diagnosis? *Online Journal of Otolaryngology and Rhinology*. 2020; 4(1). doi: 10.33552/OJOR.2020.04.000578.
- [17] Ahmed S, Ashfaq M, Shabbir A. Modified Quilting Sutures: A New Technique for Hematoma and Abscess of Nasal Septum. *Journal of College of Physicians and Surgeons Pakistan*. 2016 Jun; 26(6): 531-2.
- [18] Karodpati N, Ingale M, Rawat S, Kuradagi V. Comparing the Outcome of Septoplasty and Septoplasty with Turbinectomy in Patients with Deviated Nasal Septum. *International Journal of Otorhinolaryngology Head Neck Surg*. 2019 Sep; 5(5): 1. doi: 10.18203/issn.2454-5929.ijohns20193616.
- [19] Majumder AS, Hossen MI, Mahmud Z, Choudhury MA. Clinical Evaluation of Septoplasty with and without Additional Nasal Procedures. *International Surgery Journal*. 2025; 12(6). doi:10.18203/2349-2902.isj20251531.
- [20] Ceresoli M, Ferro CA, La Greca A, Cioffi SP, Biloslavo A, Podda M et al. Minimally Invasive Approach to Peritonitis from Left Colonic Perforation: A Retrospective Multicenter Observational Study. *Surgical Endoscopy*. 2025 Feb; 1-0. doi:10.1007/s00464-025-11611-w.