



Original Article



Evaluation of Factors Leading to the Development of Acute Pancreatitis at Tertiary Care Hospital, Karachi

Muneeba Asif^{1*}, Faizan Tariq², Faraz Tarique³, Abdul Wadood Shah⁴, Rukhsana Soomro⁵ and Aliya Zaman⁶

¹Department of Medicine, Muhammad Medical and Dental College, Ibn e Sina University, Mirpurkhas, Pakistan

²Department of Internal Medicine, Murshid Hospital and Health Care Centre, Karachi, Pakistan

³Department of Infectious Diseases, Shaheed Mohtarma Benazir Bhutto, Institute of Trauma, Karachi, Pakistan

⁴Department of Community Medicine, Bilawal Medical College, Jamshoro, Pakistan

⁵Department of Physiology, Suleman Roshan Medical College, Tando Adam, Pakistan

⁶Department of Pathology, Muhammad Medical and Dental College, Ibn e Sina University, Mirpurkhas, Pakistan

ARTICLE INFO

Keywords:

Cholelithiasis, Alcohol, Post-ERCP, Hypertriglyceridemia, IgG4-Mediated Pancreatitis

How to Cite:

Asif, M., Tariq, F., Tarique, F., Shah, A. W., Soomro, R., & Zaman, A. (2025). Evaluation of Factors Leading to the Development of Acute Pancreatitis at Tertiary Care Hospital, Karachi: Evaluation of Factors Leading to the Development of Acute Pancreatitis. *Pakistan Journal of Health Sciences*, 6(7), 287-292. <https://doi.org/10.54393/pjhs.v6i7.3347>

***Corresponding Author:**

Muneeba Asif
 Department of Medicine, Muhammad Medical and Dental College, Ibn e Sina University, Mirpurkhas, Pakistan
muneebaasif94@gmail.com

Received Date: 30th May, 2025

Revised Date: 22nd July, 2025

Acceptance Date: 28th July, 2025

Published Date: 31st July, 2025

ABSTRACT

Acute pancreatitis (AP) involves pancreatic inflammation that may extend to surrounding tissues. Clinical outcomes can be predicted through assessment of disease-specific prognostic markers. **Objectives:** To evaluate the prevalence of etiological factors for AP among patients at a tertiary healthcare center in Karachi. **Methods:** A descriptive cross-sectional design, this six-month study was conducted in the Medicine Department of Civil Hospital, Karachi, from 06-08-2022 to 05-02-2023. Participants were selected through convenience sampling (a non-probability approach). Data processing was done by SPSS version 26.0. Chi-square test was applied by taking a p-value of ≤ 0.050 , which is significant. **Results:** The research cohort comprised 135 eligible participants with a mean age of 46.71 ± 6.82 years. Gender distribution showed 55 male patients (40.7%) and 80 female patients (59.3%). Out of the 135 patients, 43.7% had cholelithiasis, 11.1% had alcohol induced pancreatitis, 17.8% had post-ERCP pancreatitis, 14.8% had hypertriglyceridemia, and 10.4% had IgG4-mediated pancreatitis. **Conclusions:** Acute pancreatitis is a multi-factorial pathology of different severity and outcomes. Gallstones were found to be the most prevalent cause of the condition.

INTRODUCTION

Acute pancreatitis (AP) is a form of inflammation of the pancreas. Although it has a mild course in most cases and does not cause serious complications in about 80% of cases, it may cause serious complications, and up to 20% may also succumb to it in some individuals [1]. A patient with acute epigastric pain that radiates to the back, nausea and vomiting must also be urgently examined to exclude several potentially fatal illnesses, among which there are

myocardial infarction, rupture or aortic aneurysm, peptic ulcer disease and its complications of hemorrhage or penetrating injury, and acute pancreatitis [2]. The patient's history should guide the clinician, who should conduct a physical inspection to assess specific issues and review the results of appropriate diagnostic tests to establish a proper treatment plan by narrowing the range of potential diagnoses [3, 4]. The prevalence of risk factors is known to

vary geographically and demographically, which is one reason the causes and incidence of acute pancreatitis differ across geographic regions [5]. Common etiological factors include alcoholic pancreatitis with higher overall acute pancreatitis incidence in certain countries as compared to countries with predominantly gallstone pancreatitis, as it is reported differently in various countries in terms of incidence curve [6]. Hospital visits in emergency departments due to acute pancreatitis have been on the rise. In 2012, the emergency visits rate for AP was 105 per 100,000 population, and the general mortality rate was approximately 1%. Death is more likely to occur when age increases, the presence of comorbidity and greater severity of diseases, especially caused by infected pancreatic necrosis and multiorgan failure [7]. Gallstones are the most prevalent causative factor of acute pancreatitis, whereas alcohol is more prominent among middle-aged men. The unknown cause known as idiopathic cases causes 20% to 34% of the cases, with no difference in their rates of occurrence among men and women [8-10]. The rate of idiopathic pancreatitis may vary depending on the clinician's diligence in seeking the underlying cause of the condition, as reported by patients [11]. Autoimmune pancreatitis (AIP) is an infrequent nonneoplastic disease, which is considered a different manifestation of pancreatic ductal adenocarcinoma (PDAC). The level of serum immunoglobulin G4 (IgG4) is useful for distinguishing between these disorders, as elevated IgG4 is a characteristic feature of AIP but not observed in PDAC patients [12]. Hameed et al., reported the etiologic distribution of acute pancreatitis (AP) in their cohort: 74% cholelithiasis, 2.6% alcohol-induced pancreatitis, 4.5% post-ERCP pancreatitis, and 0.6% hypertriglyceridemia with IgG4-mediated pancreatitis [13]. Efficient prevention measures and treatment regimens are still important interventions recommended to be adopted globally to limit the morbidity and deaths due to acute pancreatitis. Although significant progress has been made in the pathophysiological understanding of this condition over the last fifteen years, it remains challenging for clinicians to manage its multifactorial aetiology and complex clinical manifestations [14]. There are also changes in lifestyle, demographic shifts, socioeconomic status, and morbidities that continue to evolve. These factors should be acknowledged in the prevention of the onset of irreversible chronic pancreatitis, as the most valuable measures to prevent its occurrence must have been established previously [15]. Ultimately, understanding the various causes that contribute to the development of acute pancreatitis is crucial for effectively managing and preventing the condition. This study provides a benchmark that can be shared with other healthcare facilities, which

will be the study's key result. Finally, there is a need to understand the factors leading to the development of acute pancreatitis for its better management and prevention. Hence, the results from this study form the benchmark that is shared with other healthcare facilities. The findings of this study are expected to provide valuable insights into the unique clinical profile of acute pancreatitis in this population, enabling the development of targeted interventions and enhancing the overall management of this challenging condition.

Acute pancreatitis is a common gastrointestinal emergency with diverse etiological factors such as gallstones, alcohol consumption, metabolic disorders, and post-procedural complications. The distribution of these risk factors varies across different populations due to demographic, lifestyle, and healthcare differences. In Pakistan, particularly in tertiary care hospitals, limited local data are available regarding the relative prevalence of these etiological factors among patients presenting with acute pancreatitis. This lack of region-specific evidence creates a gap in understanding the dominant causes and limits the development of targeted preventive and management strategies. This study aims to evaluate the prevalence of etiological factors for AP among patients at a tertiary healthcare center in Karachi.

METHODS

A descriptive cross-sectional study was conducted at the Department of Medicine, Civil Hospital, Karachi, from 06-08-2022 to 05-02-2023 after getting ethical approval from the CPSP research evaluation unit (Ref No: CPSP/REU/MED-2019-183-15622) and written informed consent was obtained from the patients. Using the WHO sample size calculator with a 4.5% post-ERCP incidence rate [13], 3.5% margin of error, and 95% confidence interval, we calculated a required sample of 135 subjects. A design effect (DEFF) of 1 was applied to account for a simple random sampling design. The total population size was estimated at 20,000. The study utilized a non-probability consecutive sampling methodology. Inclusion criteria was confined on newly diagnosed patients with acute pancreatitis presenting within 24 hours, either gender, patients having age of 30-60 years were included and patients with history of hepatitis B, C and HIV, patients with history of malignancy, pregnant women, patients with history of asthma, COPD, stroke, congestive heart failure, Patients with chronic liver disease and chronic kidney disease were excluded from the study. The patient's history was taken in detail, and they underwent a clinical examination upon admission. The researcher also examined possible causes of acute pancreatitis development, such as cholelithiasis (diagnosed with the help of an abdominal ultrasound), alcohol abuse, post-

ERCP surgery, hypertriglyceridemia (a lipid profile), and IgG4-related pancreatitis (a sample of IgG4). A quantitative research design was employed in this study; participants' age was treated as a continuous variable. Qualitative variables were gender, residential status, type 2 diabetes mellitus, hypertension, smoking status and the risk factors of acute pancreatitis. The data analysis was conducted using the SPSS version 26.0. Means were calculated as the mean value \pm standard deviation (SD) in cases of normally distributed quantitative variables. Qualitative variables, including gender, residence status, type II diabetes mellitus, hypertension, smoking status, and all factors related to acute pancreatitis (cholelithiasis, alcohol consumption, post-ERCP, hypertriglyceridemia, and IgG4-mediated pancreatitis), were calculated in terms of frequency and percentages. To measure the modification effect of the factors, stratification was based on the attributes of age, gender, residence status, type 2 diabetes mellitus, hypertension, and smoking status. This enabled them to be evaluated regarding their effects on the outcome variables. Following stratification of the data, statistical analysis was conducted using chi-square or Fisher's exact tests, where a p-value ≤ 0.050 denoted significance.

RESULTS

This research analyzed 135 patients treated at the Civil Hospital Karachi's Department of Medicine, Pakistan, who met the study's inclusion and exclusion criteria. The participants' ages varied from 30 to 60 years, with a mean age of 46.71 ± 6.82 years. Regarding the gender distribution, 55 (40.7%) were male and 80 (59.3%) were female. Regarding the status of residence, 43 (31.9%) were residing in urban locations, while 92 (68.1%) were in rural locations. Regarding hospital stay, most of the patients stay for ≤ 5 days (66.67%) (Table 1).

Table 1: Demographic, Clinical, and Biochemical Characteristics of Patients

Variables	Frequency (%)	Mean \pm SD
Age (Years)		
30-45	103 (76.30%)	46.71 \pm 6.82
46-60	32 (23.70%)	
Gender		
Male	55 (40.70%)	-
Female	80 (59.30%)	
Residence		
Urban	43 (31.90%)	-
Rural	92 (68.10%)	
Length of Hospital Stay		
≤ 5	90 (66.67%)	-
> 5	45 (33.33%)	

Findings demonstrate risk factors causing pancreatitis.

Out of 135 patients, 59 (43.7%) and 76 (56.3%) had and did not have cholelithiasis, respectively. Out of 135 patients, 15 (11.10%) had alcohol, and 120 (88.90%) did not have respectively. Out of 135 patients, 24 (17.8%) had post-ERCP, and 111 (82.2%) did not have post-ERCP, respectively. Out of 135 patients, 20 (14.8%) had hypertriglyceridemia, and 115 (85.2%) did not have hypertriglyceridemia, respectively. Out of 135 patients, 14 (10.4%) had IgG4-mediated disease, and 121 (89.6%) did not have IgG4-mediated disease, respectively, where chi-square results indicate a p-value of 0.001 (Table 2).

Table 2: Risk Factors Causing Pancreatitis

Variables	Frequency (%)	p-Value
Cholilithiasis	59 (43.70%)	0.001
Alcohol	15 (11.10%)	0.001
Post ERCP	24 (17.80%)	0.002
Hypertriglyceridemia	20 (14.8%)	0.002
IgG4 mediated	14 (10.4%)	0.003

Results shows the associated factors causing pancreatitis. Frequency distribution on diabetes mellitus type II showed that out of 135 patients, 97 (71.90%) had diabetes mellitus type II while 38 (28.10%) did not have diabetes mellitus type II respectively. In the case of hypertension, 46 (34.1) patients were dominated by hypertension, whereas 89 (65.9) were non-hypertensive. About smoking status, 15 (11.1%) patients were smokers, while 88.9% of patients were non-smokers (Table 3).

Table 3: Associated Factors

Variables	Frequency (%)
Diabetes Mellitus Type II	
Yes	97 (71.90%)
No	38 (28.10%)
Hypertension	
Yes	46 (34.10%)
No	89 (65.90%)
Smoking	
Yes	15 (11.10%)
No	112 (88.90%)

An age and gender stratification of cholelithiasis revealed that among individuals aged 30-45 years, 45 (43.7%) had cholelithiasis, while 58 (56.3%) did not. In the 46-60 years age group, cholelithiasis was exhibited in 14 (43.8%), whereas most had no such occurrence (18, 56.2%). The relationship between cholelithiasis and age was not statistically significant, with a p-value of 0.990. In the stratification by gender, 15 (27.3%) out of the total of males had cholelithiasis, whereas 40 (72.7%) were found to possess none. Among the female patients, 44 (55%) had cholelithiasis, while 36 (45%) did not. The p-value was large ($p=0.010$), indicating a significant association (Table 4).

Table 4: Stratification for Age and Gender with Respect to Cholelithiasis

Variables	Cholelithiasis		p-Value
	Yes	No	
Age (Years)			
30-45	45 (43.7%)	58 (56.3%)	0.990
46-60	14 (43.8%)	18 (56.2%)	
Gender			
Male	15 (27.3%)	40 (72.7%)	0.010
Female	44 (55%)	36 (45%)	

DISCUSSION

Acute pancreatitis is an interesting disease, categorized by the inflammation of pancreatic tissue, parenchyma oedema, and necrosis due to auto-digestion of the glandular enzymes, which can cause multi-organ failure or even death [16]. Gallstones represent the most common etiology of acute pancreatitis in Pakistan, whereas alcohol-related cases are relatively less frequent. In daily routine practice, drug-induced acute pancreatitis is not very common [17]. Over the last several decades, the intensive care of patients with AP has improved significantly because of its connection with high morbidity and mortality rates. The aetiology of the disease must be determined to effectively handle and prevent further occurrences of the disease. Two of the most frequent etiological factors, i.e. alcohol and gallstones, account for a total of 80%, and alcoholic pancreatitis is by far much more prevalent [18]. According to the recent recommendations, the current aetiology of AP must be determined in at least 80% of cases, and not more than 20% can be seen as idiopathic. The management and administration of care to AP should be planned according to the knowledge of the equivalence as well as the aetiology of the disease [19]. According to the study done by Hameed *et al.* out of the 154 patients, 59.1% were women [13]. The most frequent aetiology encountered was gallstones (74%), followed by idiopathic (13%). A cholecystectomy performed on 44.7% of gallstone pancreatitis patients is called a same-admission cholecystectomy. Upon admission, 14.9% of patients had serum amylase levels within three times the upper normal limit, while 29 (18.8%) exhibited serum lipase levels within the same range. A greater score on the Ranson and modified CT severity index translated to a greater triage and death percentage. The rate of complications was 21.4 % with a mortality rate to be 7.1%. There were two hundred and twelve (212) patients in a research study conducted by Patel *et al.*, [20], with 88 men (73.33%) and 32 women (26.66%). The average age of the participants was 36.96 years, with a standard deviation of 13.44 years. The most frequent presenting symptom was abdominal pain, and then came vomiting. Alcohol use was the major cause of acute pancreatitis, as it was the cause in 85 patients

(70.8 per cent). The cause was identified in 25 patients (20.8%), and in five patients (4.1%), it was unknown. There were three deaths as a result of severe pancreatitis among the patients (2.5%). Additionally, a body mass index (BMI) of 25 kg/m² or higher, a hematocrit (HCT) of 44% or more, and a C-reactive protein (CRP) level of more than 150 mg/L were associated with a higher risk of developing severe acute pancreatitis. Datta *et al.*, used a sample of 40 patients with acute pancreatitis, comprising 20 male and 20 female, with an average age of 44.3 ± 2.7 years [21]. Out of the 40 cases, 26 patients (65%) received a diagnosis of moderate to severe acute pancreatitis. Certain clinical parameters, such as ascites or pleural effusion, were not identified to have significant correlations with the severity of the illness. 62.5% of the cases displayed gallstones and metabolic conditions such as hypertriglyceridemia and hypercalcemia; however, none of them could be, in any way, significantly associated with the severity of the disease.

This study was conducted at a single tertiary care center with a relatively limited sample size and utilized convenience sampling, which may affect the generalizability of the findings. Additionally, the cross-sectional design restricts the ability to establish causal relationships between risk factors and the development of acute pancreatitis. Long-term clinical outcomes and recurrence rates were not assessed. Future multicenter prospective studies with larger sample sizes and extended follow-up are recommended to better understand etiological patterns and improve preventive and therapeutic strategies for acute pancreatitis.

CONCLUSIONS

AP is still a typical diagnosis that causes hospitalization. Gallstones are the most frequent cause, and post-ERCP comes next. It has a great influence on institutional costs, hospitalization and death. The disease severity may be mild or severe and entails a risk of having systemic complications. The main points of management are timely fluid resuscitation, adequate nutrition support, and management of complications at the moment they occur. Patients with mild acute pancreatitis due to gallstones are normally recommended to undergo cholecystectomy before their release to avoid any recurrence. A multidisciplinary approach is encouraged when looking at moderate to severe cases of acute pancreatitis, especially with local and systemic complications.

Authors' Contribution

Conceptualization: MA

Methodology: MA, AWS, RS, AZ

Formal analysis: MA

Writing and Drafting: FT¹, FT², AWS, AZ

Review and Editing: FT¹, FT², AWS, AZ, MA, RS

All authors approved the final manuscript and take responsibility for the integrity of the work

Conflicts of Interest

All the authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Samad A, Ahmed A, Zehra B, Iftikhar J. Aetiology, Severity and Outcome of Patients Admitted with Acute Pancreatitis: A Cross-Sectional Study. *Journal of the Dow University of Health Sciences*. 2024 Mar; 18(1). doi: 10.36570/jduhs.2024.1.2047.
- [2] Shah AP, Mourad MM, Bramhall SR. Acute Pancreatitis: Current Perspectives on Diagnosis and Management. *Journal of Inflammation Research*. 2018 Mar; 77-85. doi: 10.2147/JIR.S135751.
- [3] Abbasi AH and Abid S. Acute Pancreatitis: A Comparative Study to Assess the Clinical Outcome of Acute Pancreatitis in Relation with Major Etiologies. *The Professional Medical Journal*. 2019 Apr; 26(04): 658-62. doi: 10.29309/TPMJ/2019.26.04.3373.
- [4] Khan ZA, Shahab F, Khan RA. Management of Acute Pancreatitis: Experience of A Tertiary Care Hospital in Peshawar. *Anesthesia, Pain and Intensive Care*. 2019 Jan; 176-9.
- [5] Shehzad MI, Shah M, Zada I, Qureshi NK, Azeem SB, Khokhar A. Clinical Outcomes of Acute Pancreatitis in Patients with Cirrhosis. *Pakistan Journal of Medical and Health Sciences*. 2023 Apr; 17(02): 672-. doi: 10.53350/pjmhs2023172672.
- [6] Iqbal F, Majeed S, Islam OS, Khan IA, Ud Din MJ, Zia MK. Frequency of Gall Stones in Patients with Acute Pancreatitis at Tertiary Care Hospital of Karachi, Pakistan: Frequency of Gall Stones in Acute Pancreatitis. *Allied Medical Research Journal*. 2024 Jan; 2(1): 213-20. doi: 10.59564/amrj/02.01/024.
- [7] Pak LM, Schattner MA, Balachandran V, D'Angelica MI, DeMatteo RP, Kingham TP et al. The Clinical Utility of Immunoglobulin G4 in the Evaluation of Autoimmune Pancreatitis and Pancreatic Adenocarcinoma. *Hepato-Pancreato-Biliary*. 2018 Feb; 20(2): 182-7. doi: 10.1016/j.hpb.2017.09.001.
- [8] Chatterjee R, Parab N, Sajjan B, Nagar VS. Comparison of Acute Physiology and Chronic Health Evaluation II, Modified Computed Tomography Severity Index, and Bedside Index for Severity in Acute Pancreatitis Score in Predicting the Severity of Acute Pancreatitis. *Indian Journal of Critical Care Medicine: Peer-Reviewed, Official Publication of Indian Society of Critical Care Medicine*. 2020 Feb; 24(2): 99. doi: 10.5005/jp-journals-10071-23343.
- [9] Kazmi SJ, Zafar MT, Zia BF, Khalid SR, Kumar V, Tabassum S et al. Role of Serum C-Reactive Protein (CRP)/Albumin Ratio in Predicting the Severity of Acute Pancreatitis: A Retrospective Cohort. *Annals of Medicine and Surgery*. 2022 Oct; 82: 104715. doi: 10.1016/j.amsu.2022.104715.
- [10] Shafiq F, Khan MF, Asghar MA, Shamim F, Sohaib M. Outcome of Patients with Acute Pancreatitis Requiring Intensive Care Admission: A Retrospective Study from A Tertiary Care Center of Pakistan. *Pakistan Journal Of Medical Sciences*. 2018 Sep; 34(5): 1082. doi: 10.12669/pjms.345.15575.
- [11] Anees F, Khan RS, Naz S, Wadani ZH. Acute Pancreatitis in Pregnancy and Puerperium: Assessing Maternal and Fetal Impact, Etiologies, and Clinical Outcomes at a Tertiary Care Hospital in Pakistan. *Cureus*. 2024 Dec; 16(12). doi: 10.7759/cureus.76393.
- [12] Akram M, Arsalan M, Kumar V, Ahmed M, Rizwan M, Iqbal W et al. Outcome of Laparoscopic Cholecystectomy in Acute Pancreatitis Secondary to Gallstones: A Prospective Observational Study from Pakistan. *Hypertension*. 2022; 170:44-.
- [13] Hameed T, Khan AZ, Shah I, Khan AQ, Ahmad M. Factors Leading to Acute Pancreatitis in Tertiary Care Hospitals in Pakistan—A Multicenter Study. *Journal of Rawalpindi Medical College*. 2018 Mar; 22(1).
- [14] Islam MS, Hossain A, Islam MD, Bala CS, Abira M, Rahman A. Clinical Profile, Etiology and In-hospital Outcome of Acute Pancreatitis: Experience at a Tertiary Care Center, Bangladesh. *Bangladesh Journal of Medicine*. 2025; 36(1): 32-6. doi: 10.3329/bjm.v36i1.77864.
- [15] Bhattarai S and Gyawali M. Clinical Profile and Outcomes in Patients with Acute Pancreatitis Attending A Teaching Hospital at Gandaki Province, Nepal. *Journal of College of Medical Sciences-Nepal*. 2020 Sep; 16(3): 168-72. doi: 10.3126/jcmsn.v16i3.31299.
- [16] Barreto SG, Habtezion A, Gukovskaya A, Lugea A, Jeon C, Yadav D et al. Critical Thresholds: Key to Unlocking the Door to the Prevention and Specific

- Treatments for Acute Pancreatitis. *Gut*. 2021 Jan; 70(1): 194–203. doi: 10.1136/gutjnl-2020-322163.
- [17] Zhou Y, Ge YT, Shi XL, Wu KY, Chen WW, Ding YB *et al*. Machine Learning Predictive Models for Acute Pancreatitis: A Systematic Review. *International Journal of Medical Informatics*. 2022 Jan; 157: 104641. doi: 10.1016/j.ijmedinf.2021.104641.
- [18] Trikudanathan G, Yazici C, Phillips AE, Forsmark CE. Diagnosis and Management of Acute Pancreatitis. *Gastroenterology*. 2024 Sep; 167(4): 673–88. doi: 10.1053/j.gastro.2024.02.052.
- [19] Konda N, Venkateswarlu^o D, Rao BT, Suresh V, Shaik A, Manasa D. Clinical Approach, Diagnosis and Medical Management of Acute. *International Journal of Research in Medical Sciences*. 2023 Jul; 11(7): 2631. doi: 10.18203/2320-6012.ijrms20232112.
- [20] Patel ML, Shyam R, Atam V, Bharti H, Sachan R, Parihar A. Clinical Profile, Etiology, and Outcome of Acute Pancreatitis: Experience at A Tertiary Care Center. *Annals of African Medicine*. 2022 Apr; 21(2): 118–23. doi: 10.4103/aam.aam_83_20.
- [21] Datta IK, Haque MN, Bhuiyan TM. Clinical Profile, Degree of Severity and Underlying Factors of Acute Pancreatitis among A Group of Bangladeshi Patients. *Ibrahim Medical College Journal of Medical Science*. 2018 Jan; 12(1): 6–10. doi: 10.3329/imcjms.v12i1.35170.